

オペレーションズ・リサーチ

3

関数 $f: \mathbb{R}^n \rightarrow \mathbb{R}$ は 2 回連続的微分可能な関数とし、 \mathbf{a} は $\mathbf{0}$ でない n 次元ベクトルとする。
次の非線形計画問題を考える。

$$\begin{aligned} P: \text{Minimize } & f(\mathbf{x}) \\ \text{subject to } & \mathbf{a}^\top \mathbf{x} = 0 \end{aligned}$$

ただし、 $^\top$ はベクトルの転置を表す。 \mathbf{x}^* を問題 P の大域的最適解とする。

さらに、次の非線形計画問題を考える。

$$\begin{aligned} P(k): \text{Minimize } & f_k(\mathbf{x}) \\ \text{subject to } & (\mathbf{x} - \mathbf{x}^*)^\top (\mathbf{x} - \mathbf{x}^*) \leq 1 \end{aligned}$$

ただし、 k は非負の整数であり、 $f_k: \mathbb{R}^n \rightarrow \mathbb{R}$ は以下に定義された関数である。

$$f_k(\mathbf{x}) = f(\mathbf{x}) + \frac{k}{2}(\mathbf{a}^\top \mathbf{x})^2 + \frac{1}{2}(\mathbf{x} - \mathbf{x}^*)^\top (\mathbf{x} - \mathbf{x}^*)$$

問題 $P(k)$ の大域的最適解を \mathbf{x}^k とする。さらに、 $\lim_{k \rightarrow \infty} \mathbf{x}^k = \bar{\mathbf{x}}$ 、 $\lim_{k \rightarrow \infty} k(\mathbf{a}^\top \mathbf{x}^k) = \bar{\lambda}$ と仮定する。

以下の問いに答えよ。

- (i) 任意の非負の整数 k に対して $f_k(\mathbf{x}^k) \leq f(\mathbf{x}^*)$ が成り立つことを示せ。
- (ii) $\mathbf{a}^\top \bar{\mathbf{x}} = 0$ 、 $\bar{\mathbf{x}} = \mathbf{x}^*$ となることを示せ。
- (iii) 問題 $P(k)$ のカルーシュ・キューン・タッカー (Karush-Kuhn-Tucker) 条件を書け。
- (iv) 十分大きな k に対して、 $\nabla f_k(\mathbf{x}^k) = \mathbf{0}$ となることを示せ。
- (v) $\nabla f(\mathbf{x}^*) + \bar{\lambda} \mathbf{a} = \mathbf{0}$ となることを示せ。

An English Translation:

Operations Research

3

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice continuously differentiable function, and let \mathbf{a} be an n -dimensional nonzero vector.

Consider the following nonlinear programming problem:

$$\begin{aligned} \text{P: Minimize } & f(\mathbf{x}) \\ \text{subject to } & \mathbf{a}^\top \mathbf{x} = 0, \end{aligned}$$

where the superscript \top denotes transposition of a vector. Let \mathbf{x}^* be a global optimal solution to problem P.

Moreover consider the following nonlinear programming problem:

$$\begin{aligned} \text{P}(k): \text{ Minimize } & f_k(\mathbf{x}) \\ \text{subject to } & (\mathbf{x} - \mathbf{x}^*)^\top (\mathbf{x} - \mathbf{x}^*) \leq 1, \end{aligned}$$

where k is a nonnegative integer and $f_k : \mathbb{R}^n \rightarrow \mathbb{R}$ is the function defined by

$$f_k(\mathbf{x}) = f(\mathbf{x}) + \frac{k}{2}(\mathbf{a}^\top \mathbf{x})^2 + \frac{1}{2}(\mathbf{x} - \mathbf{x}^*)^\top (\mathbf{x} - \mathbf{x}^*).$$

Let \mathbf{x}^k be a global optimal solution to problem P(k). Suppose that $\lim_{k \rightarrow \infty} \mathbf{x}^k = \bar{\mathbf{x}}$ and $\lim_{k \rightarrow \infty} k(\mathbf{a}^\top \mathbf{x}^k) = \bar{\lambda}$.

Answer the following questions.

- (i) Show that $f_k(\mathbf{x}^k) \leq f(\mathbf{x}^*)$ for any nonnegative integer k .
- (ii) Show that $\mathbf{a}^\top \bar{\mathbf{x}} = 0$ and $\bar{\mathbf{x}} = \mathbf{x}^*$.
- (iii) Write out Karush-Kuhn-Tucker conditions for problem P(k).
- (iv) Show that $\nabla f_k(\mathbf{x}^k) = \mathbf{0}$ for all k sufficiently large.
- (v) Show that $\nabla f(\mathbf{x}^*) + \bar{\lambda} \mathbf{a} = \mathbf{0}$.