

Exact Algorithms for Maximum Independent Set ¹

MINGYU XIAO

School of Computer Science and
Engineering, University of Electronic
Science and Technology of China, China,
myxiao@gmail.com

HIROSHI NAGAMOCHI

Department of Applied Mathematics and
Physics, Graduate School of Informatics,
Kyoto University, Japan,
nag@amp.i.kyoto-u.ac.jp

Abstract

We show that the maximum independent set problem (MIS) on an n -vertex graph can be solved in $1.1996^n n^{O(1)}$ time and polynomial space, which even is faster than Robson's $1.2109^n n^{O(1)}$ -time exponential-space algorithm published in 1986. We also obtain improved algorithms for MIS in graphs with maximum degree 6 and 7, which run in time of $1.1893^n n^{O(1)}$ and $1.1970^n n^{O(1)}$, respectively. Our algorithms are obtained by using fast algorithms for MIS in low-degree graphs in a hierarchical way and making a careful analyses on the structure of bounded-degree graphs.

Key words. Exact Algorithm, Independent Set, Graph, Polynomial-space, Branch-and-reduce, Measure-and-conquer,

1 Introduction

Over the last few decades, an extensive research has been done on exact exponential algorithms. Many interesting methods and results have been obtained in this area, which can be found in a nice survey by Woeginger [14] and a recent monograph by Fomin and Kratsch [5]. In the line of research on worst-case analysis of exact algorithms for NP-hard problems, the *maximum independent set* problem (MIS) is undoubtedly one of the most fundamental problems. The problem is used to test the efficiency of some new techniques of exact algorithms and often introduced as the first problem in some textbooks and lecture notes of exact algorithms. However, despite of a large number of contributions on exact algorithms and their worst-case analyses for MIS during the last 30 years, no published algorithm runs faster than the $1.2109^n n^{O(1)}$ -time exponential-space algorithm by Robson in 1986 [10]. Fomin and Kratsch say that ‘the running time of current branching algorithms for MIS with more and more detailed analyses seems to converge somewhere near 1.2^n ’ [5]. Researchers are interesting in how fast we can exactly solve MIS and believe that some new techniques are required to get a further significant improvement.

Related work. The first nontrivial exact algorithm for MIS is back to Tarjan and Trojanowski's $2^{n/3} n^{O(1)}$ -time algorithm in 1977 [12]. Later, Jian obtained a $1.2346^n n^{O(1)}$ -time algorithm [7]. Robson gave a $1.2278^n n^{O(1)}$ -time polynomial-space algorithm and a $1.2109^n n^{O(1)}$ -time exponential-space algorithm [10]. Robson also claimed better running times in a technical report [11]. Fomin *et al.* [4] introduced the ‘‘measure-and-conquer’’ method and got a simple $1.2210^n n^{O(1)}$ -time polynomial-space algorithm by using this method. Also based on this method, Kneis *et al.* [8] and Bourgeois *et al.* [1] improved the running time bound to $1.2132^n n^{O(1)}$ and $1.2114^n n^{O(1)}$ respectively, which are the current fastest polynomial-space algorithms for MIS in published articles. There is also a large amount of contributions to MIS in degree-bounded graphs [9, 6, 15, 16, 17, 18]. Let MIS- i mean MIS in graphs with maximum degree i . Now MIS-3 can be solved in $1.0836^n n^{O(1)}$ time [16], MIS-4 can be solved in $1.1376^n n^{O(1)}$ time [17], MIS-5 can be solved in $1.1737^n n^{O(1)}$ time [18] and MIS-6 can be solved in $1.2050^n n^{O(1)}$ time [1], where

¹ **Technical report 2013-006, November 1, 2013.** A preliminary version of this paper was presented at the 24th international symposium on algorithms and computation (ISAAC 2013).

all of them use only polynomial space. The measure-and-conquer method is a powerful tool to design or analyze exact algorithms. Most fast polynomial-space algorithms for MIS are designed based on the method. By combining this method with a bottom-up method, Bourgeois *et al.* [1] got the $1.2114^n n^{O(1)}$ -time polynomial-space algorithm for MIS. Their algorithm is based on fast algorithms for MIS in low-degree graphs.

Our contributions. In this paper, we will design a $1.1996^n n^{O(1)}$ -time polynomial-space algorithm for MIS, which is faster than Robson’s $1.2109^n n^{O(1)}$ -time exponential-space algorithm [10] obtained in 1986. We also show that MIS-6 and MIS-7 can be solved in $1.1893^n n^{O(1)}$ and $1.1970^n n^{O(1)}$ time, respectively. Our algorithms use the measure-and-conquer method. But the improvement is not obtained by studying more cases in previous algorithms. Instead, we will introduce some new methods to reduce a large number of cases and make the algorithm and its analysis easy to follow. Our algorithms also need to use our previous fast algorithms for MIS in low-degree graphs. The improvement is mainly obtained by using the following ideas:

1. We exploit a divide-and-conquer method to get the improved algorithms for MIS in high-degree graphs based on fast algorithms for MIS in low-degree graphs. In the method, we design an algorithm for MIS made of two procedures. One procedure is an algorithm to solve MIS in graphs with maximum degree *at most* i . The other procedure is to effectively deal with vertices of degree *at least* $i + 1$ in the graph. We also use the idea to design fast algorithms for MIS in degree bound graphs. Once an algorithm for MIS- i is obtained, we design a procedure for eliminating vertices of degree at least $i + 1$ by reduction/branching operations, which together with the algorithm for MIS- i will give an algorithm for MIS- $(i + 1)$. Similar bottom-up ideas have been used in some previous algorithms, such as the algorithm for MIS in [1] and the algorithm for the parameterized vertex cover problem in [2]. One advantage of our method is that, the divide-and-conquer method can combine the measure-and-conquer method well to design exact algorithms. Then we can catch the properties of fast algorithms for MIS in low-degree graphs and propagates the improvement from instances of low-degree graphs to those of high-degree graphs.
2. We devise a method that can reduce a huge number of case analyses in the algorithms and then our algorithms become much easier to check the correctness. This method is based on Lemma 6 in Section 4. It can also be directly used to reduce a large number of cases in the analysis of previous algorithms without modifying the algorithms.
3. We introduce a new branching rule, called “branching on edges,” to deal with edges between end-vertices with many common neighbors, for which the standard branching on a vertex of maximum degree has not lead to a sufficiently high performance to improve the previous time bounds.

2 Preliminaries

2.1 Notation system

Let $G = (V, E)$ stand for a simple undirected graph with a set V of vertices and a set E of edges. Let $|G|$ denote $|V|$. We will use n to denote $|V| = |G|$ and n_i to denote the number vertices of degree i in G , and $\alpha(G)$ to denote the size of a maximum independent set of G . The vertex set and edge set of a graph G are denoted by $V(G)$ and $E(G)$, respectively. For simplicity, we may denote a singleton set $\{v\}$ by v .

For a vertex subset $X \subseteq V$ in a graph G , we define the following notations. Let $G - X$ denote the graph obtained from G by removing X together with edges incident on any vertex in X , $G[X] = G - (V - X)$ be the graph induced from G by the vertices in X , and G/X denote the

graph obtained from G by contracting X into a single vertex (removing self-loops and parallel edges). Also we let $N(X)$ denote the set of all vertices in $V - X$ that are adjacent to a vertex in X , and $N[X] = X \cup N(X)$.

For a vertex v in a graph G of maximum degree d , we define the following notations. Let $\delta(v) = |N(v)|$ denote the degree of v , and $N_2(v)$ denote the set of vertices with distance exactly 2 from v , and let $N_2[v] = N_2(v) \cup N[v]$. Let e_v denote the number of edges in the subgraph $G[N(v)]$ induced by the neighbors of v (i.e., $e_v = |E(G[N(v)])|$), let f_v denote the number of edges between $N[v]$ and $N_2(v)$, and let q_v denote the number of vertices of degree $< d$ in $N_2(v)$. Also define the *neighbor-degree* k_v of v to be the sequence (k_1, k_2, \dots, k_d) (where d is the maximum degree of the graph) of the number k_i of degree- i neighbors $u \in N(v)$. Then $\sum_{1 \leq i \leq d} ik_i = \sum_{u \in N(v)} \delta(u) = \delta(v) + 2e_v + f_v$. We may denote $k_v = (k_3, k_4, k_5, k_6)$ when $k_1 = k_2 = 0$ and $k_i = 0$ for $i \geq 7$.

For each neighbor $u \in N(v)$ of v , we call a vertex $z \in N(u)$ adjacent to v (resp., not adjacent to v) the *inner-neighbor* of u at v (resp., *outer-neighbor* of u at v). Define the *inner-degree* (resp., *outer-degree*) of u at v to be the number of inner-neighbors (resp., outer-neighbors) of u at v .

2.2 Branching algorithms and the measure-and-conquer method

Our algorithms use a branch-and-reduce paradigm. We branch the current problem instance into several smaller instances to search a solution. The iterative algorithm will create a search tree. To scale the size of the instance, we need to select a measure for it. A common measure of a graph problem is the number of vertices or edges in the graph. By bounding the size of the search tree to a function of the measure, we will get a running time bound related to the measure for the problem. In MIS, a branching rule will branch on the current instance G into several instances G_1, G_2, \dots, G_l such that the measure μ_i of each G_i is less than the measure μ of G , and a solution to G can be found in polynomial time if a solution to each of the l instances G_1, G_2, \dots, G_l is known. Usually, G_i ($i = 1, 2, \dots, l$) are obtained by deleting some vertices in G . We will use $C(\mu)$ to denote the worst-case size of the search tree in the algorithm when the measure of the instance is at most μ . The above branch creates the recurrence relation $C(\mu) \leq \sum_{i=1}^l C(\mu - \mu'_i)$, where $\mu'_i = \mu - \mu_i$. The largest root of the function $f(x) = 1 - \sum_{i=1}^l x^{-\mu'_i}$, denoted by $\tau(\mu'_1, \mu'_2, \dots, \mu'_l)$, is also called the *branching factor* of the above recurrence relation. Let τ be the maximum branching factor among all branching factors in the search tree. Then the size of the search tree is $C(\mu) = O(\tau^\mu)$. More details about the analysis and how to solve recurrences can be found in the monograph [5].

In some cases, the worst branch in the algorithm will not always happen. We can use the following idea of amortization to get better analysis. Consider two branching operations A and B with recurrences $C(\mu) \leq C(\mu - t_{(A1)}) + C(\mu - t_{(A2)})$ and $C(\mu) \leq C(\mu - t_{(B1)}) + C(\mu - t_{(B2)})$ such that the branching operation B leads to a better recurrence (with a smaller branching factor) than A does, where the recurrence for the branching operation A may be the bottleneck in the run time analysis of the algorithm. Suppose that branching operation B is always applied to the subinstance G_1 generated by the first branch of A in the algorithm. In this case, we can obtain a better recurrence than that for A if we derive a recurrence by combining branching operation A and branching operation B applied to G_1 . However, in general, there may be many branching operations B_1, B_2, \dots that can be applied to G_1 . To ease such an analysis without generating all combined recurrences, we introduce a notion of “shift.” To improve the branching factor of the recurrence for operation A , we transfer some amount from the measure decrease in the recurrence for operation B to that for A as follows. We save an amount $\sigma > 0$ of measure decrease from B by evaluating the branch operation B with recurrence

$$C(\mu) \leq C(\mu - (t_{(B1)} - \sigma)) + C(\mu - (t_{(B2)} - \sigma)),$$

which is worse than its original recurrence. The saved measure decrease σ will be included into the recurrence for operation A to obtain

$$C(\mu) \leq C(\mu - (t_{(A1)} + \sigma)) + C(\mu - t_{(A2)}).$$

The saved amount is also called a *shift*, where the best value for σ will be determined so that the maximum branching factor τ is minimized. In our algorithm, we introduce one shift σ in the analysis of our algorithm for MIS-6.

To reduce the size of the search tree, we wish to find good branching rules, and try to avoid using bad branching rules with poor performance in designing algorithms. The selection of the measure is also an important issue in order to evaluate how quickly problem instances can decrease after each branching operation. The measure-and-conquer method [4] allows us to define a sophisticated way of measuring the size of problem instances. In this method, we set a weight to each vertex in the graph according to the degree of the vertex (usually vertices of the same degree receive the same weight) and define the sum of the weights in the graph to be the measure. Note that when a vertex v is deleted, we may decrease the measure not only from v but also from the neighbors of v since the degrees of the neighbors will decrease by 1. This yields an effect of amortizing branching factors from several different recurrences. Compared to the traditional measures, the weighted measure may catch more structural information of the graph and leads to a further improvement without modifying the algorithms; in fact, algorithms can be designed so that the target measure decreases as fast as possible before a final algorithm is proposed. Currently, the best exact algorithms for many NP-hard problems are designed by using this method. An important step in this method is to set vertex weights as valuables. We sometime solve a quasiconvex program to determine the best values of them to minimize the maximum branching factor τ . In this paper we also employ the branch-and-reduce paradigm as our algorithms and the measure-and-conquer method to analyze their run times.

2.3 Reduction operations

Before applying our branching rules, we may first apply some reduction rules to reduce some local structures, branching on which may lead to a bad performance. Reduction rules can be applied in polynomial time to find a part of the solution or decrease the size of the instance directly. Many nice reduction rules have been developed. In this paper, we only use three known reduction rules.

Reduction by removing unconfined vertices

A vertex v in an instance G is called *removable* if $\alpha(G) = \alpha(G - v)$. A sufficient condition for a vertex to be removable has been studied in [16]. In this paper, we only use a simple case of the condition. A neighbor $u \in N(v)$ of v is called an *extending child* of v if u has exactly one outer-neighbor $s_u \in N_2(v)$ at v , where s_u is also called an *extending grandchild* of v . Let $N^*(v)$ denote the set of all extending children $u \in N(v)$ of v , and S_v be the set of all extending grandchildren s_u ($u \in N^*(v)$) of v together with v itself. We call v *unconfined* if there is a neighbor $u \in N(v)$ which has no outer-neighbor or $S_v - \{v\}$ is not an independent set (i.e., some two vertices in $S_v \cap N_2(v)$ are adjacent)². It is known in [16] that any unconfined vertex is removable.

Lemma 1 [16] *For an unconfined vertex v in graph G , it holds that*

$$\alpha(G) = \alpha(G - v).$$

²Unconfined vertices in [16] are defined in a more general way.

A vertex u *dominates* another vertex v if $N[u] \subseteq N[v]$, where v is called *dominated*. We see that dominated vertices are unconfined vertices.

Reduction by folding complete k -independent sets

We call a set $A = \{v_1, \dots, v_k\}$ of k degree- $(k+1)$ vertices a *complete k -independent set* if they have common neighbors $N(v_1) = \dots = N(v_k)$.

Lemma 2 [16] *For a complete k -independent set A , we have that*

$$\alpha(G) = \alpha(G^*) + k,$$

where $G^* = G/N[A]$ if $N(A)$ is an independent set and $G^* = G - N[A]$ otherwise.

Folding a complete k -independent set A is to eliminate the set $N[A]$ from an instance in the above way. In our algorithm, we only fold complete k -independent set with $k \leq 2$, since this operation is good enough for our analysis. Folding a complete 1-independent set $A = \{v\}$ consisting of a degree-2 vertex v is also called *folding a degree-2 vertex v* .

Reduction by removing line graphs

If a graph H is the line graph of a graph H' , then a maximum independent set of H can be obtained as the set of vertices that corresponds the set of edges in a maximum matching in H' . To reduce some worst cases, we need to remove the line graphs of 4-regular graphs, the line graphs of (4,5)-bipartite graphs (a bipartite graph with edges between two sets V_1 and V_2 is a (d_1, d_2) -bipartite graph if every vertex in V_i is of degree d_i ($i = 1, 2$)) and the line graphs of 5-regular graphs. A graph is the line graph of a 4-regular graph (resp., 5-regular graph) if and only if the graph has only degree-6 vertices (resp., degree-8 vertices) and each of them is contained in two edge-disjoint cliques of size 4 (resp., 5). A graph is the line graph of a (4,5)-bipartite graph if and only if the graph has only degree-7 vertices and each of them is contained in two edge-disjoint cliques of size 4 and 5, respectively. More characterizations of line graphs can be found in [13]. Removing line graphs of 4-regular graphs (resp., (4,5)-bipartite graphs and 5-regular graphs) is useful in the analysis of our algorithm for MIS-6 (resp., MIS-7 and MIS-8).

Definition 3 *A graph is called a reduced graph, if it contains none of unconfined vertices, complete k -independent sets with $k = 1$ or 2, and a component of a line graph of a 4-regular graph, a (4,5)-bipartite graph or a 5-regular graph.*

The algorithm in Figure 1 is a collection of all above reduction operations. When the graph is not a reduced graph, we can use the algorithm in Figure 1 as a preprocessing to reduce it. Notice that even if a graph of maximum degree θ is given as an instance to MIS- θ ($\theta \in \{6, 7, 8\}$), a vertex of degree $d \geq \theta + 1$ may be created by contraction of vertices during an execution of algorithm **reduce**.

3 Divide-and-conquer method

We exploit a divide-and-conquer approach to design algorithms for solving MIS and MIS- i ($i \geq 3$). In this method, we divide the class of instances of MIS into two classes, one consisting of instances of maximum degree at least i for some $i \geq 3$, and the other consisting of those of maximum degree at most $i - 1$. For the first class of instances, we design a procedure that applies reduction/branching operations until the maximum degree of the instance decreases to at most $i - 1$. Then we switch to an algorithm that solves the second class of instances, i.e., MIS- $(i - 1)$. We combine a procedure for the instances of maximum degree at least i with an algorithm for solving MIS- $(i - 1)$ to obtain an algorithm for MIS.

Input: A graph $G = (V, E)$.

Output: A reduced graph $G' = (V', E')$ and the size $s = |S|$ of a subset $S \subseteq V - N[V']$ such that for any maximum independent set X of G' , the union $X \cup S$ is a maximum independent set to G .

Initialize $s := 0$ and $G' := G$;

Execute the following steps as long as at least one of them is applicable before returning the resulting pair (G', s) ;

1. For each component H of G' that is the line graph of a 4-regular graph, a (4,5)-bipartite graph or a 5-regular graph, compute $\alpha(H)$, and let $G' := G' - V(H)$ and $s := s + \alpha(H)$;
2. For each unconfined vertex $v \in V$, let $G' = G' - v$;
3. For each complete k -independent set A with $k = 1$ or 2 , let $G' := G^*$ and $s := s + k$ for $G^* = G - N[A]$ if $N(v)$ is an independent set, and $G^* = G/N[A]$ otherwise.

Figure 1: Algorithm $\text{reduce}(G, s)$

For MIS- j for each $3 \leq j \leq i - 1$, we can use the same idea. Namely, we split the class of instances of MIS- j into two classes, one consisting of instances of maximum degree j , the other consisting of those of maximum degree at most $j - 1$. For the first class of instances, we design a procedure that applies reduction/branching operations until the maximum degree of the instance decreases by at least one. We obtain an algorithm for MIS- j by combining a procedure for the instances of maximum degree at least j with an algorithm for solving MIS- $(j - 1)$. For MIS-3, we only need to design a procedure that applies reduction/branching operations until no vertex of degree at least 3 is left, since a maximum independent set in an instance of maximum degree at most 2 can be found in linear time.

Our algorithm for each classes of instances are designed by branch-and-paradigm and analyzed by the measure-and-conquer method. For MIS or MIS- θ , we let $w_i \geq 0$ denote the weight of a degree- i vertex in an instance of the problem such that and define the *measure* μ of a graph G with n_i degree- i vertices ($i \geq 0$) to be

$$\mu(G) = \sum_i w_i n_i.$$

We may assign different values to w_i depending on the problem MIS or MIS- θ under the condition that $w_i \leq 1$ for each vertex of a given instance of MIS or MIS- θ so that $\mu(G) \leq n$ holds (during an execution of our algorithm to MIS- θ , a vertex of degree $i > \theta$ may be created and $w_i > 1$ is allowed for such a vertex). In any of MIS and MIS- θ , we see that any instance G with $\mu(G) = 0$ can be solved in polynomial time. Hence if the measure never increases after any reduction or branching operation, then we can bound the size of search trees from above by a function $\tau^{\mu(G)}$ of $\mu(G)$ ($\leq n$).

However, sometimes it is not easy to analyze the running time of the combined algorithms since that a different measure (vertex weight) may be used for the algorithm to each class. We will introduce a method to effectively deal with this difficulty, especially for the case where the measure is set as the sum of total weight of vertices in the graph.

Let A_i denote an algorithm that solves MIS- i in a graph G of maximum degree $\leq i$ in $(\tau_i)^{\mu_i(G)} |G|^{O(1)}$ time, where τ_i is a positive number and $\mu_i(G) = \sum_{1 \leq j \leq i} w_j^{(i)} n_j$ is the measure of

G (recall that n_j is the number of degree- j vertices in G and $w_j^{(i)} \geq 0$ is the weight of a degree- j vertex). Let $B_{>i}$ denote a procedure that branches on a graph G of maximum degree $> i$ with branching factor τ'_i on measure $\mu_{i+1}(G) = \sum_{j \geq 1} w_j^{(i+1)} n_j$, where $w_j^{(i+1)} \geq 0$ is the weight of a degree- j vertex in the procedure. We have the following lemma for analyzing combined algorithms for MIS:

Lemma 4 *For an integer $i \geq 3$, let $\lambda = \max\{\frac{w_j^{(i)}}{w_j^{(i+1)}} \mid 0 \leq j \leq i, w_j^{(i+1)} \neq 0\}$ and $\tau_{i+1} = \max\{\tau'_i, (\tau_i)^\lambda\}$. Then MIS can be solved in $(\tau_{i+1})^{\mu_{i+1}(G)} |G|^{O(1)}$ time.*

Proof. We will construct an algorithm A_{i+1} that solves MIS in $O^*((\tau_{i+1})^{\mu_{i+1}(G)})$ time. It iteratively applies the procedure $B_{>i}$ to branch when the graph has maximum degree $> i$, and calls the algorithm A_i when the graph has maximum degree at most i . We analyze the running time of A_{i+1} .

In A_{i+1} , we use $\mu_{i+1}(G)$ as the measure (the same measure in $B_{>i}$). When the graph has a vertex of degree at least $i+1$, the algorithm can branch with branching factor $\tau'_i \leq \tau_{i+1}$. When the graph becomes a graph of maximum degree at most i , the algorithm will execute A_i . In this part, the algorithm uses time $O^*((\tau_i)^{\mu_i(G_0)})$, where $\mu_i(G_0) = \sum_{1 \leq j \leq i} w_j^{(i)} n_j \leq \lambda \sum_{1 \leq j \leq i} w_j^{(i+1)} n_j = \lambda \mu_{i+1}(G_0)$ (note that $n_j = 0$ for $j > i$). This implies that the algorithm can always branch with branching factor $(\tau_i)^\lambda \leq \tau_{i+1}$ on measure $\mu_{i+1}(G)$ in this part. Therefore, the algorithm A_{i+1} runs in $O^*((\tau_{i+1})^{\mu_{i+1}(G)})$ time. \blacksquare

We can analyze combined algorithms for MIS- θ .

Here is an application of Lemma 4. We will show that MIS-8 can be solved in time $1.19951^{\mu_8(G)} |G|^{O(1)}$ time, where $\mu_8(G) = 0.65844n_3 + 0.78844n_4 + 0.88027n_5 + 0.95345n_6 + 0.98839n_7 + n_8$, and that in a graph with maximum degree at least 9 we can branch with branching factor 1.19749 on the measure $\mu_9(G) = \sum_j n_j$. In Lemma 4, we have $\tau'_8 = 1.19749$, $\tau_8 = 1.19951$, and $\lambda = \max\{0.65844, 0.78844, 0.88027, 0.95345, 0.98839, 1\} = 1$. Then MIS can be solved in $1.19951^n |G|^{O(1)}$ time.

In the above method, we let $\tau_{i+1} = \max\{\tau'_i, (\tau_i)^\lambda\}$, where τ'_i is decided by $B_{>i}$, τ_i is decided by A_i , and λ is related to the vertex weights in both of $B_{>i}$ and A_i . So sometimes simple reductions on τ'_i or τ_i may not lead to improvement on the algorithm A_{i+1} . To get more properties and further improvements on the problem, in our algorithm, we may not design A_i and $B_{>i}$ totally independently. Instead, we will design $B_{>i}$ based on A_i by considering the result (the values of τ_i and vertex weight) of A_i as some constraints to set the vertex weight in $B_{>i}$.

This divide-and-conquer method provides a way to solve MIS by solving two subproblems and to design fast algorithms for MIS based on fast algorithms for MIS in low-degree graphs. We will focus on the subalgorithm $B_{>i}$. Fast algorithms A_i for MIS- i with $i = 3, 4$ and 5 can be found in references [16, 17, 18].

In this paper, by using this divide-and-conquer method, first, we design an algorithm for MIS-6 based on fast algorithm for MIS-5 in [18], second, we design an algorithm for MIS-7 based on the algorithm for MIS-6, third, we design an algorithm for MIS-8 based on the algorithm for MIS-7, and finally, we design an algorithm for MIS in general graphs based on the algorithm for MIS-8. Our results are listed in Table 1.

4 Branching on High-Degree Vertices

There is an easy way to deal with high-degree vertices. We can simply branch on a high-degree vertex v into two branches by including it to the solution set or not. In the branch where v is included to the solution, $N[v]$ will be deleted from the graph since the neighbors of v cannot be

Problem	Running time	Vertex weight	References
MIS-6	$1.18922^n n^{O(1)}$	$(w_3, w_4, w_5) = (0.49969, 0.76163, 0.92401)$	Section 7
MIS-7	$1.19698^n n^{O(1)}$	$(w_3, w_4, w_5, w_6) = (0.65077, 0.78229, 0.89060, 0.96384)$	Section 8
MIS-8 (MIS)	$1.19951^n n^{O(1)}$	$(w_3, w_4, w_5, w_6, w_7) =$ $(0.65844, 0.78844, 0.88027, 0.95345, 0.98839)$	Section 9

Table 1: Our algorithms designed by the divide-and-conquer method

selected into the solution anymore. If the degree of v is higher, then the graph can be reduced more in this branch. We extend the simple branch rule based on this following observation. For a vertex v , there are only two possible cases: (i) there is a maximum independent set of the graph which does not contain v ; and (ii) every maximum independent set of the graph contains v . Recall here the set S_v of all extending grandchildren of v together with v itself. As is shown in [16], we see that for Case (ii), S_v is always contained in any maximum independent set of the graph. We get the following branching rule.

Branching on a vertex v means generating two subinstances by excluding v from the independent set or including S_v to the independent set. In the first branch we will delete v from the instance whereas in the second branch we will delete $N[S_v]$ from the instance.

Branching on a vertex v of maximum degree d is one of the most fundamental operations in our algorithm. We analyze this operation. Throughout the paper, we use $\Delta w_i = w_i - w_{i-1}$ ($i \geq 3$), and assume that

$$0 \leq \Delta_{i+1} \leq \Delta_i \quad (i \geq 2); \quad 2\Delta_\theta \leq \Delta_{\theta-1} \text{ for MIS-}\theta \ (\theta \in \{6, 7, 8\}) \quad (1)$$

(these inequalities will be automatically satisfied with the optimized weights w_i in our algorithms to MIS- θ).

Let $\Delta_{out}(v)$ and $\Delta_{in}(v)$ to denote the decrease of the measure of μ in the branches of excluding v and including S_v , respectively. Then we get recurrence $C(\mu) = C(\mu - \Delta_{out}(v)) + C(\mu - \Delta_{in}(v))$. By letting $k_v = (k_3, k_4, \dots, k_d)$ be the neighbor-degree of v , we give more details about lower bounds on $\Delta_{out}(v)$ and $\Delta_{in}(v)$.

For the first branch, we get

$$\Delta_{out}(v) = w_d + \sum_{i=3}^d k_i \Delta w_i.$$

Observe that, for a fixed neighbor-degree k_v , the decrease $\Delta_{out}(v)$ in the first branch is small when the neighbors of v have higher degrees since $\Delta_i \geq \Delta_{i+1}$.

In the second branch, we will delete $N[S_v]$ from the graph. Let $\Delta(\overline{N[v]})$ denote the decrease of weight of vertices in $V(G) - N[v]$ by removing $N[S_v]$ from G together with possibly weight decrease attained by reduction operations applied to $G - N[S_v]$. Then we have

$$\Delta_{in}(v) \geq w_d + \sum_{i=3}^d k_i w_i + \Delta(\overline{N[v]}).$$

We observe that, for a fixed neighbor-degree k_v , the decrease $\Delta_{in}(v)$ in the second branch is determined by $\Delta(\overline{N[v]})$.

Then we can branch on a vertex v of maximum degree d with recurrence

$$\begin{aligned} C(\mu) &= C(\mu - \Delta_{out}(v)) + C(\mu - \Delta_{in}(v)) \\ &\leq C(\mu - (w_d + \sum_{i=3}^d k_i \Delta w_i)) + C(\mu - (w_d + \sum_{i=3}^d k_i w_i + \Delta(\overline{N[v]}))). \end{aligned} \quad (2)$$

A simple lower bound on $\Delta(\overline{N[v]})$ is obtained as follows.

Lemma 5 For a vertex v of maximum degree d , it holds

$$\Delta(\overline{N[v]}) \geq (f_v + (f_v - |N_2(v)|) + q_v)\Delta w_d \geq f_v\Delta w_d.$$

Proof. Let ℓ_z denote the number of edges between $N(v)$ and z , where $f_v = \sum_{z \in N_2(v)} \ell_z$. Since the degree of z decreases by ℓ_z after removing $N[v]$ from G , we see that $\Delta(\overline{N[v]}) \geq \sum_{z \in N_2(v)} (w_{\delta(z)} - w_{\delta(z) - \ell_z})$. The number of degree- d vertices in $N_2(v)$ is $|N_2(v)| - q_v$ (recall that q_v is the number of vertices of degree $\leq d-1$ in $N_2(v)$). Since $\Delta_i \geq \Delta_{i+1}$ and $\Delta_{d-1} \geq 2\Delta_d$, we have $\sum_{z \in N_2(v)} (w_{\delta(z)} - w_{\delta(z) - \ell_z}) = \sum_{z \in N_2(v): d(z)=d} (w_d - w_{d-\ell_z}) + \sum_{z \in N_2(v): d(z)<d} (w_{d-1} - w_{d-1-\ell_z}) \geq (|N_2(v)| - q_v)\Delta w_d + q_v\Delta w_{d-1} + \Delta w_{d-1} \sum_{z \in N_2(v)} (\ell_z - 1) \geq (|N_2(v)| - q_v)\Delta w_d + 2\Delta w_d q_v + 2\Delta w_d (f_v - |N_2(v)|) = (f_v + (f_v - |N_2(v)|) + q_v)\Delta w_d$, as required. \blacksquare

We here remark about some feature on the recurrence (2). In the recurrence (2), usually the measure decrease in the first branch of removing a vertex v is much smaller than that in the second branch, and the branching factor of the recurrence tends to be easily large when the measure decrease in the first branch is small; i.e., the neighbors of v have higher degrees. Another remark is a special effect of the condition of $N^*(v) \neq \emptyset$ to the term $\Delta(\overline{N[v]})$. Recall that the second branch of including S_v into the solution removes not only $N[v]$ but also $N[S_v] - N[v]$. This provides a larger lower bound on $\Delta(\overline{N[v]})$ than that in Lemma 5 (see Lemma 15(ii) for a detailed analysis).

As for branching on vertices of maximum degree in our algorithms, we examine the recurrence (2) for all possible neighbor-degrees (k_3, k_4, \dots, k_d) to evaluate the branching factor precisely, and show the existence of vertices that attain a large value in the lower bound on $\Delta(\overline{N[v]})$ in Lemma 5 based on a graph theoretical argument.

Before closing this section, we propose a new method for knowing the maximum branching factor of recurrences (2) over all neighbor-degrees of v . Assume that we use a fixed lower bound on $\Delta(\overline{N[v]})$. A straightforward method is to create a concrete recurrence for each neighbor-degree $k_v = (k_3, k_4, \dots, k_d)$ of v . However, the number of neighbor-degrees $k_v = (k_3, k_4, \dots, k_d)$ is $(d+1)^{d-3}$ (the number of integer solutions to the function $k_3 + k_4 + \dots + k_d = d$). Thus (2) actually consists of $(d+1)^{d-3}$ concrete recurrences. We introduce a technical lemma that can eliminate redundant recurrences to determine the largest branching factor among a set of systematically generated recurrences. With this, we can reduce the number of recurrences in (2) from $(d+1)^{d-3}$ to only $d-2$.

Lemma 6 Let $C(x) = \tau^x$ for a positive $\tau > 1$. For any nonnegative $p, \mu, a_i, b_i, i = 1, 2, \dots, \ell$ ($\ell \geq 1$), the maximum of

$$C(\mu - (\sum_{i=1,2,\dots,\ell} k_i a_i + c)) + C(\mu - (\sum_{i=1,2,\dots,\ell} k_i b_i + d))$$

over all $k_1, k_2, \dots, k_\ell \geq 0$ subject to $k_1 + k_2 + \dots + k_\ell = p$ is equal to the maximum of

$$C(\mu - (pa_i + c)) + C(\mu - (pb_i + d))$$

over all $i = 1, 2, \dots, \ell$.

Proof. It suffices to show that for nonnegative $w, a_1, b_1, b_2, c, d \geq 0$, it holds $C(\mu - (a_1 + a_2 + c)) + C(\mu - (b_1 + b_2 + d)) \leq \max\{C(\mu - (2a_1 + c)) + C(\mu - (2b_1 + d)), C(\mu - (2a_2 + c)) + C(\mu - (2b_2 + d))\}$. The lemma can be obtained by applying this repeatedly. Note that function $f(t) = \tau^{-(2a_1(1-t) + 2a_2t + c - \mu)} + \tau^{-(2b_1(1-t) + 2b_2t + d - \mu)}$ is convex since the second derivative is nonnegative. Hence $f(0.5) \leq \max\{f(0), f(1)\}$ holds, as required. \blacksquare

By applying Lemma 6, in (2), we only need to consider $d - 2$ concrete recurrences with neighbor-degrees $(k_3, k_4, \dots, k_d) = (d, 0, \dots, 0), (0, d, 0, \dots, 0), \dots, (0, \dots, 0, d)$, respectively. For example, if $d = 6$, we can decrease the number of recurrences from $7^4 = 2401$ to only 5. Lemma 6 is introduced to simplify the analysis of recurrences for the first time. It can be used to reduced thousands of recurrences in the analysis of previous algorithms for MIS, such as the algorithms in [8] and [1]. Note that the authors of [8] used a computer-added method to create all possible recurrences in the web page [20]. There are more than 10 thousands recurrences listed. By using Lemma 6, we need to generate a set of about 50 recurrences, which is now easily checkable by hand.

5 Branching on Edges

As we have remarked in the previous section, the maximum branching factor of recurrences (2) becomes larger when $N^*(v) = \emptyset$, v has high degree neighbors and $\Delta(\overline{N[v]})$ is small. By considering that $\Delta(\overline{N[v]}) = f_v \Delta_d$ can hold in Lemma 5, we wish to avoid branching on a vertex v with a small f_v , particularly when $N^*(v) = \emptyset$ and $k_d = d$.

Our solution to this situation is to introduce a new branching rule that can deal with the dense local graph $G[N(v)]$ caused for a small f_v (a large e_v). That is “branching on edges.”

Branching on edges Two disjoint independent subsets A and B of vertices in a graph G are called *alternative* if $|A| = |B| \geq 1$ and there is a maximum independent set S_G of G which satisfies $S_G \cap (A \cup B) = A$ or B . Let G^\dagger be the graph obtained from G by removing $A \cup B \cup (N(A) \cap N(B))$ and adding an edge ab for every two nonadjacent vertices $a \in N(A) - N[B]$ and $b \in N(B) - N[A]$.

Lemma 7 [16] *For alternative subsets A and B in a graph G , $\alpha(G) = \alpha(G^\dagger) + |A|$.*

Lemma 8 *Let vv' be an edge. Then*

$$\alpha(G) = \max\{\alpha(G - \{v, v'\}), \alpha(G^\dagger) + 1\},$$

where G^\dagger be the graph obtained from G by removing $\{v, v'\} \cup (N(v) \cap N(v'))$ and adding an edge ab for every two nonadjacent neighbors $a \in N(v) - N[v']$ and $b \in N(v') - N[v]$.

Proof. We easily observe that either (i) every maximum independent set S_G of G satisfies $S_G \cap \{v, v'\} = \emptyset$; or (ii) there is a maximum independent set S_G of G such that $S_G \cap \{v, v'\} \neq \emptyset$. In (i), we have $\alpha(G) = \alpha(G - \{v, v'\})$. In (ii), sets $A = \{v\}$ and $B = \{v'\}$ are alternative in G , and we have $\alpha(G) = \alpha(G^\dagger) + 1$ by Lemma 7. ■

Branching on an edge vv' means generating two subinstances according to Lemma 8. This is to either remove $\{v, v'\}$ from the graph G or construct the graph G^\dagger from $G - (\{v, v'\} \cup (N(v) \cap N(v')))$ by making each pair $a \in N(v) - N[v']$ and $b \in N(v') - N[v]$ adjacent. Branching on an edge may not always be very effective. In our algorithms, we will apply it to edges vv' when $N(v) \cap N(v')$ is large, which are called “short edges.”

We denote our algorithm for solving an instance of MIS- θ by $\text{mis}\theta(G)$. The definitions of “short edges” in algorithm $\text{mis}\theta(G)$ are slightly different with θ . In $\text{mis}\theta(G)$, an edge vv' in a reduced graph of maximum degree $\theta \in \{6, 7, 8\}$ is called *short* if

- (i) $\delta(v) = 6$, $\delta(v') \in \{5, 6\}$ and $|N(v) \cap N(v')| \geq 3$ for $\theta = 6$; and
- (ii) $\delta(v) = \delta(v') = \theta$ and $|N(v) \cap N(v')| \geq 4$ for $\theta \in \{7, 8\}$.

A short edge is called *optimal* if $|N(v) \cap N(v')| - \delta(v')$ is maximized. In our algorithms, we only branch on optimal short edges in graphs of maximum degree 6, 7 and 8.

6 Algorithms

In this section, we describe our algorithms $\text{mis}\theta(G)$, $\theta = 6, 7, 8$, and then discuss MIS in general graphs.

6.1 Algorithms for MIS-6, MIS-7 and MIS-8

Our algorithm $\text{mis}\theta(G)$, $\theta \in \{6, 7, 8\}$ is simple:

First keep branching on vertices of maximum degree $d > \theta$, then keep branching on short edges, choosing an optimal one, and then keep branching on vertices of maximum degree θ , choosing an “optimal” one. During the execution, we switch to an algorithm for MIS- $(\theta - 1)$ whenever the maximum degree of the graph becomes smaller than θ .

See Figure 2 for their descriptions. In the rest of this section, we describe which vertices of maximum degree should be chosen as “optimal” vertices. When no short edge is left in a graph G of maximum degree $\theta \in \{6, 7, 8\}$, the inner-degree of each neighbor u of a degree θ -vertex v is at most 2 for $\theta = 6$ and 3 for $\theta \in \{7, 8\}$ (otherwise vu would be a short edge). In particular, for every degree θ -vertex v with $N^*(v) = \emptyset$, the outer-degree of every neighbor $u \in N(v)$ is at least 2 at v , and we have $f_v \geq 2\delta(v) + k_{\delta(v)}$ for $\theta \in \{6, 7\}$; $f_v \geq 2\delta(v) + 2k_{\delta(v)}$ for $\theta = 8$. As we have remarked, we know that the branching factor of recurrence (2) tends to be larger when $N^*(v)$ is empty and the neighbor-degree k_v consists of higher degrees. As a vertex v to branch on first should be one with $N^*(v) \neq \emptyset$, or with a neighbor-degree k_v which is lexicographically small. When k_v is close to $(k_3 = 0, 0, \dots, 0, k_\theta = \theta)$, we choose a vertex that attains a large value in the lower bound $(f_v + (f_v - |N_2(v)|) + q_v)$ in Lemma 5. This is our priority for selecting vertices of maximum degree θ . We define “optimal” vertices for each $\text{mis}\theta(G)$, $\theta = 6, 7, 8$ according to this.

In a reduced graph of maximum degree 6 in MIS-6, a degree-6 vertex v is called *optimal* if at least one of the following (i)-(vi) is holds:

- (i) $k_3 \geq 1$ or $k_6 \leq 3$;
- (ii) $k_6 = 4$ and $k_5 \leq 1$;
- (iii) $k_6 = 4$, $k_5 = 2$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 17$;
- (iv) $k_6 = 5$, $k_4 = 1$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 18$;
- (v) $k_6 = 5$, $k_5 = 1$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 19$; and
- (vi) $k_6 = 6$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 22$.

In a reduced graph of maximum degree 7 in MIS-7, a degree-7 vertex v is called *optimal* if at least one of the following (i)-(iv) is holds:

- (i) $N^*(v) \neq \emptyset$;
- (ii) the vertex v has at most 5 degree-7 neighbors ($k_7 \leq 5$);
- (iii) $k_7 = 6$ and $f_v + (f_v - |N_2(v)|) \geq 22 - 2k_3 - k_4$; and
- (iv) $k_7 = 7$ and $f_v + (f_v - |N_2(v)|) \geq 26$.

In a reduced graph of maximum degree 8 in MIS-8, a degree-8 vertex v is called *optimal* if (i) $k_8 \leq 7$ or (ii) $k_8 = 8$ and $f_v + (f_v - |N_2(v)|) \geq 36$.

Note that in the definitions of optimal vertices in graphs of maximum degree 7 and 8, we do not need to use q_v .

Lemma 9 *Let G be a reduced graph of maximum degree θ in MIS- θ ($\theta \in \{6, 7, 8\}$). If G has no short edges, then G has at least one optimal vertex.*

In order to focus on the mechanism of our algorithms first, we move the proof of this purely analytical lemma to Section 10.

Input: A graph G .

Output: The size of a maximum independent set in G .

1. Reduce the graph by $(G, s) := \text{reduce}(G, 0)$, and let d be the maximum degree of G .
2. **If** $\{d \geq (\theta + 1)\}$, pick up a vertex v of maximum degree d , and **return** $s + \max\{\text{mis}\theta(G - v), |S_v| + \text{mis}\theta(G - N[S_v])\}$.
3. **Elseif** $\{d = \theta$ and G has a short edge $\}$, pick up an optimal short edge vv' , and **return** $s + \max\{\text{mis}\theta(G - \{v, v'\}), 1 + \text{mis}\theta(G^\dagger)\}$.
4. **Elseif** $\{d = \theta$ (G has no short edges) $\}$, pick up an optimal degree- θ vertex v , and **return** $s + \max\{\text{mis}\theta(G - v), |S_v| + \text{mis}\theta(G - N[S_v])\}$.
5. **Else** $\{\text{The maximum degree of } G \text{ is smaller than } \theta\}$, use our algorithm for MIS- $(\theta - 1)$ to solve the instance G and **return** $s + \alpha(G)$, where the algorithm for MIS-5 is in [18].

Note: With a few modifications, the algorithm can deliver a maximum independent set.

Figure 2: Algorithms $\text{mis}\theta(G)$

In our algorithms $\text{mis}\theta$ ($\theta = 6, 7, 8$), we set the vertex weight w_i ($i \geq 3$) as follows (recall that $w_1 = w_2 = 0$): For $3 \leq i \leq \theta - 1$, w_i is set as that in Table 1; and

$$w_\theta = 1; \quad w_i = w_\theta + (i - \theta)\Delta w_\theta \quad i \geq \theta + 1. \quad (3)$$

To simplify our analyses, the weights of vertices of degree $\geq \theta + 1$ are allowed to be greater than 1. Recall that a vertex of degree $\geq \theta + 1$ may be created after applying reduction rules. As will be observed, we create vertices of degree $\geq \theta + 1$ only when the measure does not increase. This ensures that the running time bound of our algorithms still can be expressed by $\tau^n n^{O(1)}$ with the largest branching factor τ .

Lemma 10 *With the above vertex weight setting, each recurrence generated by the algorithm $\text{mis6}(G)$ (resp., $\text{mis7}(G)$ and $\text{mis8}(G)$) in Figure 2 has a branching factor not greater than 1.18922 (resp., 1.19698 and 1.19951).*

We will give detailed analysis of our algorithms $\text{mis6}(G)$, $\text{mis7}(G)$ and $\text{mis8}(G)$ in Sections 7, 8 and 9, respectively, and then complete a proof of Lemma 10. Since the measure μ is not greater than the number n of vertices in the initial graph in $\text{mis6}(G)$, $\text{mis7}(G)$ and $\text{mis8}(G)$, we establish the next.

Theorem 11 *A maximum independent set in an n -vertex graph with maximum degree $\theta \in \{6, 7, 8\}$ can be found in time of $1.1893^n n^{O(1)}$ for $\theta = 6$, $1.1970^n n^{O(1)}$ for $\theta = 7$, and $1.1996^n n^{O(1)}$ for $\theta = 8$, respectively.*

6.2 MIS in general graphs

Our algorithm for MIS in general graphs is also simple. It only contains two steps: Keep branching on a vertex of maximum degree while the degree of the graph is 9 or larger, and invoke

our algorithm `mis8` for MIS-8 whenever the maximum degree of the graph becomes less than 9. For the procedure for dealing with vertices of degree ≥ 9 , we set the measure as the number of vertices in the graph (the weight of each vertex is 1). Then we can get the following recurrence:

$$C(\mu) \leq C(\mu - 1) + C(\mu - 10), \quad (4)$$

which has a branching factor 1.19749, better than 1.19951 for MIS-8. The analysis in Section 3 shows that MIS in general graphs can also be solved in $1.19951^n n^{O(1)}$ time.

Theorem 12 *A maximum independent set in an n -vertex graph can be found in $1.1996^n n^{O(1)}$ time and polynomial space.*

7 Analysis of `mis6`(G)

For MIS-6, we first give some properties of the subgraph $G[N(v)]$ of the neighbors of an optimal vertex v , and then derive recurrences for all branching operations in `mis6`(G).

7.1 Weight setting

Recall that, for MIS-6, we assume that $w_0 = w_1 = w_2 = 0 \leq w_3 \leq w_4 \leq w_5 \leq w_6 = 1 \leq w_7 \leq w_8 \leq \dots$, and the values of w_3 , w_4 and w_5 will be determined after we analyze how the measure changes after each step of the algorithm.

To simplify our analysis, we assume that

$$6\Delta w_6 \leq w_3, \quad (5)$$

where this condition is satisfied by the optimized values in Table 1. We impose the next constraint in order to ensure that contracting vertices will never increase the measure.

$$w_i + w_j \geq w_{i+j-2}, \quad 3 \leq i, j \leq 6. \quad (6)$$

Lemma 13 *$w_i + w_j \geq w_{i+j-2}$ holds for all $i, j \geq 1$.*

Proof. If i or j is at most 2, say $i \leq 2$, then $w_i + w_j = w_j \geq w_{i+j-2}$. Let $i, j \geq 3$. For $3 \leq i, j \leq 5$, we have $w_i + w_j \geq w_{i+j-2}$ by (6). Let at least one of i and j , say i , is at least 6. Then we have that $i + j - 2 \geq 7$ and $w_{i+j-2} = w_i + (j - 2)\Delta w_6$ by the definition of w_k ($k \geq 7$). Since $w_j \geq (j - 2)\Delta w_6$, this implies that $w_{i+j-2} = w_i + (j - 2)\Delta w_6 \leq w_i + w_j$. ■

Lemma 14 *The measure μ of a graph G never increases in $RG(G, s)$ (Step 1 of `mis6`(G)). Moreover, in a graph of maximum degree d and minimum degree ≥ 3 , the measure μ decreases by at least $2\Delta w_d$ after applying $RG(G, s)$ if the maximum degree of the graph decreases by at least one.*

Proof. $RG(G, s)$ contains three reduction steps. In Step 1, when a component H of the line graph of a 4-regular graph, a (4,5)-bipartite graph or a 5-regular graph is removed, the measure never increases. If a vertex of maximum degree is removed in this step, then the measure decreases by at least $w_d \geq 2\Delta w_d$. In Step 2, when an unconfined vertex is removed, the measure will not increase, since the vertex weight is monotonic with the degree of the vertices. If a vertex of maximum degree is reduced, then the measure decreases either by $w_d \geq 2\Delta w_d$ from this vertex or by $2\Delta w_d$ from this vertex and a neighbor of it. In Step 3, folding a complete k -independent set ($k = 1$ or 2) is applied. By Lemma 13, we know that the measure will never increase in this step. Note that in this step, either $N[A]$ is removed or a graph $G/N[A]$ is generated by

contracting $N[A]$. For the former case, the measure decreases by at least $w_3 \geq 2\Delta w_d$. For the latter case, we can see that either the measure decreases by at least $2\Delta w_d$ directly or the new created vertex has a degree not less than the degree of any vertex in $N[A]$ and all vertices other than $N[A]$ keep the same degree, which means that the maximum degree of the graph does not decrease. This completes the proof. \blacksquare

Next we will analyze the recurrence of each branching step of $\text{mis6}(G)$.

7.2 Branching on vertices of maximum degree in Step 2 of $\text{mis6}(G)$

We will derive recurrences for branchings in Step 2 of $\text{mis6}(G)$. Let G be a reduced graph after Step 1 of $\text{mis6}(G)$. The next property holds for every vertex v in G .

Lemma 15 *Let v be a vertex in a reduced graph G , and f_v denote the number of edges between $N(v)$ and $N_2(v)$, where $f_v \geq \delta(v)$.*

- (i) *If $N^*(v) = \emptyset$, then $f_v \geq 2\delta(v)$ and $\Delta(\overline{N[v]}) \geq 2\delta(v)\Delta w_6$.*
- (ii) *If $N^*(v) \neq \emptyset$, then $\Delta(\overline{N[v]}) \geq \min\{2w_3, w_3 + 2(\delta(v) - 3)\Delta w_6\}$.*

Proof. In general, we have $\Delta(\overline{N[v]}) \geq f_v\Delta w_6$ since $\Delta w_i \geq \Delta w_6$ by (1) and $6\Delta w_6 \leq w_3$ by (5). Each neighbor of v has a neighbor in $N_2(v)$ and $f_v \geq \delta(v)$, since otherwise it would dominate some other neighbor of v .

- (i) If $N^*(v) = \emptyset$, then there are at least $2\delta(v)$ edges between $N(v)$ and $N_2(v)$ and then $f_v \geq 2\delta(v)$.
- (ii) Assume that $N^*(v) \neq \emptyset$; i.e., $S_v - \{v\} \neq \emptyset$. If $|S_v - \{v\}| \geq 2$, then clearly $\Delta(\overline{N[v]}) \geq \sum_{t \in (S_v - \{v\})} w_{\delta(t)} \geq 2w_3$. Assume that $|S_v - \{v\}| = 1$ and u is the unique vertex in $S_v - \{v\}$. Each vertex in $N^*(v)$ is adjacent to u and each vertex in $N(v) - N^*(v)$ is adjacent to at least two vertices in $N_2(v)$. Let $d^* = |N^*(v)|$. Then $\Delta(\overline{N[v]}) \geq w_{\delta(u)} + 2(\delta(v) - d^*)\Delta w_6$. Note that $\delta(u) \geq \max\{3, d^*\}$. We know that $\Delta(\overline{N[v]}) \geq \min\{w_i + 2(\delta(v) - i)\Delta w_6 \mid 3 \leq i \leq \delta(v)\} \geq w_3 + 2(\delta(v) - 3)\Delta w_6$ (by (1)). \blacksquare

In particular, for vertices v with $\delta(v) \geq 7$, we obtain $\Delta(\overline{N[v]}) \geq \min\{2\delta(v)\Delta w_6, 2w_3, w_3 + 2(\delta(v) - 3)\} \geq 12\Delta w_6$ by (5) and (1).

Now we derive recurrence of branching on a vertex of degree ≥ 7 in Step 2 of $\text{mis6}(G)$. To evaluate the largest branching factor of the recurrences (2) with the lower bound $\Delta(\overline{N[v]})$ for all $d \geq 7$, we only need to consider those for vertices with no neighbors of degree $d \geq 7$, since $\Delta w_6 = \min\{\Delta w_i\}$ for all $i \geq 3$ and $w_{i+1} \geq w_i$ for all $i \geq 3$. Furthermore, this means that we only have to consider the case of $d = 7$ in cases of $k_i = 7$ ($3 \leq i \leq 7$) by Lemma 6. Thus we get recurrences:

$$\begin{aligned} C(\mu) &\leq C(\mu - (w_d + \sum_{i=3}^d k_i \Delta w_i)) + C(\mu - (w_d + \sum_{i=3}^d k_i w_i + \Delta(\overline{N[v]}))) \\ &\leq \max_{3 \leq i \leq 7} [C(\mu - (w_7 + 7\Delta w_i)) + C(\mu - (w_7 + 7w_i + 12\Delta w_6))]. \end{aligned}$$

These recurrences will not leads to the largest branching factor. In fact, two of the recurrences for branching on short edges in Step 3 of algorithm $\text{mis6}(G)$ ((10) with $i = 6$ and $j = 6$ and (12) where $i = 6$ and $j = 6$) will be the worst recurrences. Since we know that a vertex of degree ≥ 7 always will be created after the second branch in such short edge branching, we here save a shift $\sigma > 0$ from the recurrence for branching on vertices of degree ≥ 7 so that the shift $\sigma > 0$ will be included into the recurrences for the short edge branchings. Then in this step we use the following recurrences indeed:

$$C(\mu) \leq \max_{3 \leq i \leq 7} [C(\mu - (w_7 + 7\Delta w_i - \sigma)) + C(\mu - (w_7 + 7w_i + 12\Delta w_6 - \sigma))]. \quad (7)$$

7.3 Branching on short edges in Step 3 of $\text{mis6}(G)$

We derive recurrences for branching on optimal short edges vv' in Step 3 of $\text{mis6}(G)$. Let v be a degree-6 vertex, $d' = \delta(v') \in \{5, 6\}$, and $k \in \{3, 4\}$ be the number of common neighbors of v and v' . Denote $N(v) - \{v'\} = \{u_i \mid i = 1, 2, 3, 4, 5\}$, $N(v') - \{v\} = \{u'_i \mid i = 1, 2, \dots, d' - 1\}$, where $u_i = u'_i$, $1 \leq i \leq k$ are the common neighbors, and let i^* denote the number of degree-3 vertices $u \in \{u_1, \dots, u_k\}$, where we assume that for each $i \leq i^*$, u_i is a degree-3 neighbor of v . Let $X = \{v, v'\} \cup (N(v) \cap N(v'))$. We distinguish three cases: (i) $d' = 6$ and $k = 4$; (ii) $d' = 5$ and $k = 3$; and (iii) $d' = 6$ and $k = 3$. By the optimality of the selected short edge vv' in this step, we know that: when vv' satisfies (ii) then no short edge satisfies (i); and when vv' satisfies (iii) then no short edge satisfies (i) or (ii). This is the reason why we need to define optimal short edges.

Case (i) $d' = 6$ and $k = 4$: The first branch of deleting vertices v and v' decreases the degree of u_i ($1 \leq i \leq 4$) by two and that of u_5 and u'_5 by one. Each degree-3 neighbor u_i ($i \leq i^*$) will be a degree-1 vertex in $G - \{v, v'\}$ and its unique neighbor $z_i \in N_2(v) \cap N_2(v')$ will be removed since it is an unconfined vertex, where $z_i \notin \{u_5, u'_5\}$ (otherwise u_i would dominate z_i ($= u_5, u'_5$)) and $z_\ell \neq z_{\ell'}$ for $1 \leq \ell < \ell' \leq i^*$ (otherwise $(u_\ell, u_{\ell'})$ would be a complete k -independent set which must have been reduced in **reduce**). Hence in the first branch the measure decreases by at least $2w_6 + \sum_{1 \leq i \leq 4} (w_{\delta(u_i)} - w_{\delta(u_i)-2}) + \Delta w_{\delta(u_5)} + \Delta w_{\delta(u'_5)} + i^* w_3$, where $i^* w_3$ is from deleting z_i .

In the second branch we delete $X = \{v, v', u_1, u_2, u_3, u_4\}$ to construct graph G^\dagger , joining u_5 and u'_5 with a new edge if they are not adjacent in G . Note that G has at least four edges between X and $V - X$ other than edges vu_5 and $v'u'_5$. The second branch decreases the weight of vertices in $V - (X \cup \{u_5, u'_5\})$ by at least $4\Delta w_6$ even after an edge $u_5 u'_5$ is introduced (since edges vu_5 and $v'u'_5$ are not included to evaluate the measure decrease). Hence in the second branch, the measure decreases by at least $2w_6 + \sum_{1 \leq i \leq 4} w_{\delta(u_i)} + 4\Delta w_6$. By Lemma 6, we only need to consider the following four recurrences each of which corresponds to the case of $\delta(u_1) = \delta(u_2) = \delta(u_3) \in \{3, 4, 5, 6\}$:

$$\begin{aligned} C(\mu) \leq & C(\mu - (2w_6 + 4(w_i - w_{i-2}) + 2\Delta w_6)) \\ & + C(\mu - (2w_6 + 4w_i + 4\Delta w_6)) \quad (i = 4, 5, 6); \end{aligned} \quad (8)$$

and

$$C(\mu) \leq C(\mu - (2w_6 + 8w_3 + 2\Delta w_6)) + C(\mu - (2w_6 + 4w_3 + 4\Delta w_6)). \quad (9)$$

Next, we assume that there is no short edges vv' with $\delta(v) = \delta(v') = 6$ and $|N(v) \cap N(v')| = 4$. Then the outer-degree of every degree-6 neighbor of a degree-6 vertex is at least 2.

Case (ii) $d' = 5$ and $k = 3$: Analogously with Case (i), the first branch decreases the measure by at least $w_6 + w_5 + \sum_{i=1,2,3} (w_{\delta(u_i)} - w_{\delta(u_i)-2}) + \Delta w_{\delta(u_4)} + \Delta w_{\delta(u_5)} + \Delta w_{\delta(u'_4)} + i^* w_3 \geq w_6 + w_5 + \sum_{i=1,2,3} (w_{\delta(u_i)} - w_{\delta(u_i)-2}) + 2\Delta w_6 + \Delta w_j + i^* w_3$, where $j = \delta(u'_4) \in \{3, 4, 5, 6\}$.

We consider the second branch of deleting $X = \{v, v', u_1, u_2, u_3\}$ from G to construct G^\dagger by adding edges $u_4 u'_4$ and $u_5 u'_4$, if necessary. Let p be the number of degree-6 vertices in $\{u_1, u_2, u_3\}$, where each degree-6 neighbor of v has outer-degree at least 2. Let L denote the number of edges in G between X and $V - X$ other than the three edges vu_4 , vu_5 and $v'u'_4$, where $L \geq 3 + p$. Recall that j is the degree of u'_4 in G . Then the degree of u'_4 in $G - X$ is $j - \ell - 1$, where ℓ is the number of edges between u'_4 and $\{u_1, u_2, u_3\}$. Then the degree of u'_4 in G^\dagger will be at most $j - \ell + 1$, and the weight change at vertex u'_4 from G to G^\dagger is at least

$$w_j - w_{j-\ell+1} = -(w_{j+1} - w_j) + (w_{j+1} - w_{j-\ell+1}) (\geq -(w_{j+1} - w_j) + \ell \Delta w_6).$$

Recall that $L \geq \ell$. Hence the decrease of the measure in the second branch is at least $w_6 + w_5 + \sum_{i=1,2,3} w_{\delta(u_i)} + L\Delta w_6 - (w_{j+1} - w_j) \geq w_6 + w_5 + \sum_{i=1,2,3} w_{\delta(u_i)} + (3 + p)\Delta w_6 - (w_{j+1} - w_j)$.

By Lemma 6, we only need to consider the following recurrences:

$$C(\mu) \leq C(\mu - (w_6 + w_5 + 3(w_i - w_{i-2}) + 2\Delta w_6 + \Delta w_j)) + C(\mu - (w_6 + w_5 + 3w_i + (3+p)\Delta w_6 - (w_{j+1} - w_j))), \quad (10)$$

where $4 \leq i \leq 6$ ($p = 4$ for $i = 6$ and $p = 0$ for $i = 4$ or 5) and $3 \leq j \leq 6$; and

$$C(\mu) \leq C(\mu - (w_6 + w_5 + 6w_3 + 2\Delta w_6 + \Delta w_j)) + C(\mu - (w_6 + w_5 + 3w_3 + 3\Delta w_6 - (w_{j+1} - w_j))) \quad (3 \leq j \leq 6). \quad (11)$$

We analyze a special case in (10) where $i = 6$ and $j = 6$. For this case, we get the recurrence

$$C(\mu) \leq C(\mu - (w_6 + w_5 + 3(w_6 - w_4) + 3\Delta w_6)) + C(\mu - (4w_6 + w_5 + 5\Delta w_6)).$$

In the second branch we get the graph G^\dagger , where u'_4 is a degree-7 vertex. In the next step, either the degree-7 vertex is reduced by applying a reduction rule in Step 1 or the algorithm will branch on the degree-7 vertex in Step 2. For the former case, the measure will decrease by at least $2\Delta w_6$ by Lemma 14. For the latter case, we can get σ saved from the recurrence (7). We assume that $2\Delta w_6 \geq \sigma$. Then we can get following recurrence instead of the above one

$$C(\mu) \leq C(\mu - (w_6 + w_5 + 3(w_6 - w_4) + 3\Delta w_6)) + C(\mu - (4w_6 + w_5 + 5\Delta w_6 + \sigma)).$$

Next, we further assume that there is no short edge vv' with $\delta(v) = 6$, $\delta(v') = 5$ and $|N(v) \cap N(v')| = 3$. Then the outer-degree of every degree-5 neighbor of a degree-6 vertex is at least 2.

Case (iii) $d' = 6$ and $k = 3$: Analogously with Case (ii), the first branch decreases the measure by at least $2w_6 + \sum_{i=1,2,3} (w_{\delta(u_i)} - w_{\delta(u_i)-2}) + \sum_{x \in \{u_4, u_5, u'_4, u'_5\}} \Delta w_{\delta(x)} + i^* w_3$. We consider the second branch of deleting $X = \{v, v', u_1, u_2, u_3\}$ from G to construct G^\dagger by adding edges $u_4 u'_4$, $u_4 u'_5$, $u_5 u'_4$ and $u_5 u'_5$, if necessary. Let p be the number of degree-5, 6 vertices in $\{u_1, u_2, u_3\}$, where each degree-5 or 6 neighbor of v has outer-degree at least 2. Let L denote the number of edges in G between X and $V - X$ other than the four edges vu_4 , vu_5 , $v'u'_4$ and $v'u'_5$, where $L \geq 3+p$. For each vertex $x \in \{u_4, u_5, u'_4, u'_5\}$, the degree of the vertex x in $G - X$ is $\delta(x) - \ell_x - 1$, where ℓ_x is the number of edges between x and $\{u_1, u_2, u_3\}$. Then the weight change at x from G to G^\dagger is at least

$$w_{\delta(x)} - w_{\delta(x) - \ell_x + 1} \geq -(w_{\delta(x)+1} - w_{\delta(x)}) + \ell_x \Delta w_6,$$

where $\ell_x \Delta w_6$ is the lower bound on the weight decrease caused by the deletion of the ℓ_x edges between x and $\{u_1, u_2, u_3\}$. Recall that $L \geq \sum_{x \in \{u_4, u_5, u'_4, u'_5\}} \ell_x$. Hence the measure decrease in the second branch is at least $2w_6 + \sum_{i=1,2,3} w_{\delta(u_i)} + L\Delta w_6 - \sum_{x \in \{u_4, u_5, u'_4, u'_5\}} (w_{\delta(x)+1} - w_{\delta(x)}) \geq 2w_6 + \sum_{i=1,2,3} w_{\delta(u_i)} + (3+p)\Delta w_6 - \sum_{x \in \{u_4, u_5, u'_4, u'_5\}} (w_{\delta(x)+1} - w_{\delta(x)})$. By Lemma 6, we only need to consider the following recurrences:

$$C(\mu) \leq C(\mu - (2w_6 + 3(w_i - w_{i-2}) + 4\Delta w_j)) + C(\mu - (2w_6 + 3w_i + (3+p)\Delta w_6 - 4(w_{j+1} - w_j))), \quad (12)$$

where $4 \leq i \leq 6$ ($p = 3$ for $i = 5$ or 6 ; and $p = 0$ for $i = 4$) and $3 \leq j \leq 6$; and

$$C(\mu) \leq C(\mu - (2w_6 + 6w_3 + 4\Delta w_j)) + C(\mu - (2w_6 + 3w_3 + 3\Delta w_6 - 4(w_{j+1} - w_j))), \quad (13)$$

where $3 \leq j \leq 6$.

We also analyze a special case in (12) where $i = 6$ and $j = 6$. In the second branch we get the graph G^\dagger with four degree-7 vertices u_4, u_5, u'_4 and u'_5 . Analogously with the special case in

Case (ii), in the second branch, either the measure decreases by $2\Delta w_6 \geq \sigma$ directly or we get shift σ saved from (7) by branching on a degree-7 vertex. Then for this case we can get the following recurrence instead

$$C(\mu) \leq C(\mu - (2w_6 + 3(w_6 - w_4) + 4\Delta w_6)) + C(\mu - (5w_6 + 2\Delta w_6 + \sigma)).$$

From now on, we can assume that there is no short edges vv' with $\delta(v) = \delta(v') = 6$ and $|N(v) \cap N(v')| = 3$. Then the outer-degree of every degree-6 neighbor of a degree-6 vertex is at least 3.

After Step 3 of $\text{mis6}(G)$, for each degree-6 vertex v in G , its degree-5 (resp., degree-6) neighbor is of outer-degree ≥ 2 (resp., ≥ 3) at v , and it holds

$$f_v \geq k_3 + k_4 + 2k_5 + 3k_6. \quad (14)$$

7.4 Branching on vertices of maximum degree 6 in Step 4 of $\text{mis6}(G)$

We will derive recurrences for branchings in Step 4 of $\text{mis6}(G)$. After Step 3, we can assume that the current graph G is a reduced graph with maximum degree 6 such that there is no short edge. Let v be an optimal vertex v of degree 6 selected in Step 4 of $\text{mis6}(G)$.

We define

$$\lambda_6(k_3, k_4, k_5, k_6) = \begin{cases} \min\{(12 + k_6)\Delta w_6, w_3 + 6\Delta w_6\} & \text{if } k_6 \leq 3 \text{ and } k_3 + k_4 \geq 2 \\ (6 + k_5 + 2k_6)\Delta w_6 & \text{if } k_6 \leq 3 \text{ and } k_3 + k_4 \leq 1 \\ (6 + k_5 + 2k_6)\Delta w_6 & \text{if } k_6 = 4 \text{ and } k_3 + k_4 \geq 1 \\ 17\Delta w_6 & \text{if } k_6 = 4 \text{ and } k_5 = 2 \\ (16 + 2k_4 + 3k_5)\Delta w_6 & \text{if } k_6 = 5 \\ 22\Delta w_6 & \text{if } k_6 = 6. \end{cases} \quad (15)$$

Then we have:

Lemma 16 *Let v be an optimal degree-6 vertex in Step 4 of $\text{mis6}(G)$. Then $\Delta(\overline{N[v]}) \geq \lambda_6(k_3, k_4, k_5, k_6)$.*

Proof. By (14), we have $\Delta(\overline{N[v]}) \geq f_v \Delta w_6 \geq (k_3 + k_4 + 2k_5 + 3k_6)\Delta w_6 = (6 + k_5 + 2k_6)\Delta w_6$. This proves the cases of “ $k_6 \leq 3$ and $k_3 + k_4 \leq 1$,” “ $k_6 = 4$ and $k_3 + k_4 \geq 1$,” and “ $k_6 = 5$ and $k_3 = 1$ ” (the case of “ $k_6 = 5$ and $k_3 \neq 1$ ” will be treated next).

By Lemma 5 and the definition of optimal vertices imply the case of “ $k_6 = 4$ and $k_5 = 2$,” “ $k_6 = 5$ and $k_3 \neq 1$ ” and “ $k_6 = 6$.”

Finally we show the case of “ $k_6 \leq 3$ and $k_3 + k_4 \geq 2$.” If $N^*(v) = \emptyset$ then each degree-3,4 neighbor of v also has at least two neighbors in $N_2(v)$, and we again obtain $\Delta(\overline{N[v]}) \geq f_v \Delta w_6 \geq (12 + k_6)\Delta w_6$. If $N^*(v) \neq \emptyset$, then Lemma 15(ii) implies that $\Delta(\overline{N[v]}) \geq \min\{2w_3, w_3 + 2(\delta(v) - 3)\Delta w_6\} \geq w_3 + 6\Delta w_6$. This proves the case of “ $k_6 \leq 3$ and $k_3 + k_4 \geq 2$.” ■

By Lemma 16, we obtain the recurrence (2) for $d = 6$ as follows:

$$C(\mu) \leq C(\mu - (w_6 + k_3 w_3 + k_4(w_4 - w_3) + k_5(w_5 - w_4) + k_6(w_6 - w_5))) + C(\mu - (w_6 + k_3 w_3 + k_4 w_4 + k_5 w_5 + k_6 w_6 + \lambda_6(k_3, k_4, k_5, k_6))) \quad (16)$$

for all nonnegative integers (k_3, k_4, k_5, k_6) with $k_3 + k_4 + k_5 + k_6 = 6$.

7.5 Final step

We have derived recurrences for all branching operations in algorithm `mis6`(G) except for Step 5 which invokes the fast algorithms for MIS-5 in [18]. To determine the largest branching factor to algorithm `mis6`(G) using our divide-and-conquer method in Section 3, we combine all the above recurrences with the weight setting used to determine the branching factor to algorithms for MIS-5 in [18].

The algorithm for MIS-5 in [18] runs in $1.17366^{\mu_5(G)} \mu_5(G)^{O(1)}$ time for a degree-5 graph G with measure $\mu_5(G) = \sum_{1 \leq i \leq 5} w_i^{(5)} n_i$ where n_i is the number of degree- i vertices in G , and $w_i^{(5)}$ is the weight of a degree- i vertex such that $w_0^{(5)} = w_1^{(5)} = w_2^{(5)} = 0$, $w_3^{(5)} = 0.50907$, $w_4^{(5)} = 0.82427$ and $w_5^{(5)} = 1$. Based on Lemma 4, we include the following three constraints into the current set of recurrences.

$$\begin{aligned} C(\mu) &\leq 1.17366^{\frac{w_3^{(5)}}{w_3} \mu}, & C(\mu) &\leq 1.17366^{\frac{w_4^{(5)}}{w_4} \mu}, \quad \text{and} \\ C(\mu) &\leq 1.17366^{\frac{w_5^{(5)}}{w_5} \mu}. \end{aligned} \tag{17}$$

Recurrences (7) to (13) and (16) together with (17) generate the constraints in a quasiconvex program to minimize the largest branching factor τ . By solving the quasiconvex program according to the method introduced in [3], we get an upper bound 1.18922 on the branching factor for all recurrences by setting vertex weights as

$$w_i = \begin{cases} 0 & \text{for } i = 0, 1 \text{ and } 2 \\ 0.49969 & \text{for } i = 3 \\ 0.76163 & \text{for } i = 4 \\ 0.92401 & \text{for } i = 5 \\ 1 & \text{for } i = 6 \\ w_6 + (i - 6)(w_6 - w_5) & \text{for } i \geq 7. \end{cases} \tag{18}$$

Now a feasible value of the shift σ is 0.10647. This verifies Lemma 10 with $\theta = 6$.

8 Analysis of `mis7`(G)

In the same manner of Section 7, we analyze the largest branching factor of recurrences for the branchings in `mis7`(G). All notations except for a new vertex weight in `mis7`(G) stand for the same meaning in Section 7.

Recall that, for MIS-7, we assume that $w_0 = w_1 = w_2 = 0 \leq w_3 \leq w_4 \leq w_5 \leq w_6 \leq w_7 = 1 \leq w_8 \leq \dots$, and the values of w_3 , w_4 , w_5 and w_6 will be determined after we analyze how the measure changes after each step of the algorithm.

To simplify our analysis, we assume that

$$18\Delta w_7 \leq w_3. \tag{19}$$

We impose the next constraint so that the measure does not increase after contracting vertices.

$$w_i + w_j \geq w_{i+j-2}, \quad 3 \leq i, j \leq 7. \tag{20}$$

We can see that Lemma 13 still holds in `mis7`(G) and then the measure of the graph will not increase after Step 1 of `mis7`(G). Next we derive a recurrence of each branching step of `mis7`(G).

Step 1: It is easy to see that the statement of Lemma 15 still holds for $\theta = 7$ even after replacing ‘ Δw_6 ’ with ‘ Δw_7 ’ in it. Based on the $\theta = 7$ version of Lemma 15, we see that every vertex v with $\delta(v) \geq 8$ satisfies

$$\Delta(\overline{N[v]}) \geq \min\{2\delta(v)\Delta w_7, 2w_3, w_3 + 2(\delta(v) - 3)\} \geq 14\Delta w_7$$

by (1) and (19).

Step 2: We use recurrences (2) for branching on a vertex of degree ≥ 8 in Step 2 of $\text{mis7}(G)$. By Lemma 6, we only have to consider the case of $d = 8$ for the recurrences with $k_i = 8$ ($3 \leq i \leq 8$). Thus we get recurrences

$$\begin{aligned} C(\mu) &\leq C(\mu - (w_d + \sum_{i=3}^d k_i \Delta w_i)) + C(\mu - (w_d + \sum_{i=3}^d k_i w_i + \Delta(\overline{N[v]}))) \\ &\leq \max_{3 \leq i \leq 8} [C(\mu - (w_8 + 8\Delta w_i)) + C(\mu - (w_8 + 8w_i + 14\Delta w_7))]. \end{aligned} \quad (21)$$

Step 3: We consider branching on an optimal short edge vv' in Step 3 of $\text{mis7}(G)$. We see that $|N(v) \cap N(v')| \geq 6$ cannot occur otherwise v would dominate v' .

(i) We derive recurrences in case of $|N(v) \cap N(v')| = 5$. Analogously with Case (i) in Section 7.3, we get recurrences

$$\begin{aligned} C(\mu) &\leq C(\mu - (2w_7 + 5(w_i - w_{i-2}) + 2\Delta w_7)) \\ &\quad + C(\mu - (2w_7 + 5w_i + 4\Delta w_7)) \quad (i = 4, 5, 6, 7); \end{aligned} \quad (22)$$

and

$$C(\mu) \leq C(\mu - (2w_7 + 10w_3 + 2\Delta w_7)) + C(\mu - (2w_7 + 5w_3 + 4\Delta w_7)). \quad (23)$$

(ii) We derive recurrences in case of $|N(v) \cap N(v')| = 4$. Denote $X = N(v) \cap N(v') = \{u_i \mid i = 1, 2, 3, 4\}$, $N(v) - N[v'] = \{u_5, u_6\}$ and $N(v') - N[v] = \{u'_5, u'_6\}$. Let i^* denote the number of degree-3 vertices $u \in \{u_1, u_2, u_3, u_4\}$, where we assume that for each $i \leq i^*$, u_i is a degree-3 neighbor of v . Analogously with Case (iii) in Section 7.3, the first branch of deleting vertices v and v' decreases the measure by at least $2w_7 + \sum_{i=1,2,3,4} (w_{\delta(u_i)} - w_{\delta(u_i)-2}) + \sum_{x \in \{u_5, u_6, u'_5, u'_6\}} \Delta w_{\delta(x)} + i^* w_3$. In the second branch of deleting $X = \{v, v', u_1, u_2, u_3, u_4\}$ from G to construct G^\dagger by adding edges $u_5 u'_5$, $u_5 u'_6$, $u_6 u'_5$ and $u_6 u'_6$, if necessary. Let p be the number of degree-7 vertices in $\{u_1, u_2, u_3, u_4\}$, where each degree-7 neighbor of v has outer-degree at least 2 (otherwise there would be a short edge aa' with $|N(a) \cap N(a')| \geq 5$). Let L denote the number of edges in G between X and $V - X$ other than the four edges vu_5 , vu_6 , $v'u'_5$ and $v'u'_6$, where $L \geq 4 + p$. The following analysis is the same as Case (iii) in Section 7.3. The decrease of the measure in the second branch is at least $2w_7 + \sum_{i=1,2,3,4} w_{\delta(u_i)} + (4+p)\Delta w_6 - \sum_{x \in \{u_5, u_6, u'_5, u'_6\}} (w_{\delta(x)+1} - w_{\delta(x)})$. By Lemma 6, we only need to consider the following recurrences:

$$\begin{aligned} C(\mu) &\leq C(\mu - (2w_7 + 4(w_i - w_{i-2}) + 4\Delta w_j)) \\ &\quad + C(\mu - (2w_7 + 4w_i + (4+p)\Delta w_7 - 4(w_{j+1} - w_j))), \end{aligned} \quad (24)$$

where $4 \leq i \leq 7$ ($p = 4$ for $i = 7$; and $p = 0$ for $i = 4, 5$ and 6) and $3 \leq j \leq 7$; and

$$C(\mu) \leq C(\mu - (2w_7 + 8w_3 + 4\Delta w_j)) + C(\mu - (2w_7 + 4w_3 + 4\Delta w_7 - 4(w_{j+1} - w_j))), \quad (25)$$

where $3 \leq j \leq 7$.

Step 4: We again use recurrences (2) for branching on an optimal vertex v in Step 4 of $\text{mis7}(G)$. Now the graph has no short edge, and every degree-7 vertex has outer-degree at least 3 at a degree-7 neighbor of it. Hence it holds

$$f_v \geq k_3 + k_4 + k_5 + k_6 + 3k_7. \quad (26)$$

We define

$$\lambda_7(k_7) = \begin{cases} (14+k_7)\Delta w_7 & \text{if } k_7 \leq 5 \\ (22-2k_3-k_4)\Delta w_7 & \text{if } k_7 = 6 \\ 26\Delta w_7 & \text{if } k_7 = 7. \end{cases} \quad (27)$$

Then we have:

Lemma 17 *Let v be an optimal degree-7 vertex in Step 4 of $\text{mis7}(G)$. Then $\Delta(\overline{N[v]}) \geq \lambda_7(k_7)$.*

Proof. First consider the case of $N^*(v) = \emptyset$. Then each neighbor of v has at least two neighbors in $N_2(v)$ and each degree-7 neighbor of v has at least three neighbors in $N_2(v)$. By (26), we obtain $\Delta(\overline{N[v]}) \geq f_v \Delta w_7 \geq (14+k_7)\Delta w_7$ for $k_7 \leq 5$. By Lemma 5, it holds $\Delta(\overline{N[v]}) \geq (f_v + (f_v - |N_2(v)|))\Delta w_7$. For $k_7 = 6$ (resp., $k_7 = 7$), this and the definition of optimal vertices imply that $\Delta(\overline{N[v]}) \geq (f_v + (f_v - |N_2(v)|))\Delta w_7 \geq (22-2k_3-k_4)\Delta w_7$ (resp., $\geq 26\Delta w_7$).

If $N^*(v) \neq \emptyset$, then the $\theta = 7$ version of Lemma 15(ii) implies that $\Delta(\overline{N[v]}) \geq \min\{2w_3, w_3 + 2(\delta(v) - 3)\Delta w_7\} \geq w_3 + 8\Delta w_7$, which is larger than any of $(14+k_7)\Delta w_7$, $(22-2k_3-k_4)\Delta w_7$ and $26\Delta w_7$ by (19). This proves all the cases. \blacksquare

By Lemma 16, we obtain the recurrence (2) for $d = 7$ as follows.

Case 1 ($k_7 \leq 5$):

$$C(\mu) \leq C(\mu - (w_7 + 7(w_i - w_{i-1}))) + C(\mu - (w_7 + 7w_i + 14\Delta w_7)), \quad (28)$$

where $3 \leq i \leq 6$.

Case 2 ($k_7 = 6$ and $k_3 = 1$):

$$C(\mu) \leq C(\mu - (w_7 + 6(w_7 - w_6) + w_3)) + C(\mu - (7w_7 + w_3 + 20\Delta w_7)). \quad (29)$$

Case 3 ($k_7 = 6$ and $k_4 = 1$):

$$C(\mu) \leq C(\mu - (w_7 + 6(w_7 - w_6) + (w_4 - w_3))) + C(\mu - (7w_7 + w_4 + 21\Delta w_7)). \quad (30)$$

Case 4 ($k_7 = 6$ and $k_5 + k_6 = 1$):

$$C(\mu) \leq C(\mu - (w_7 + 6(w_7 - w_6) + (w_i - w_{i-1}))) + C(\mu - (7w_7 + w_i + 22\Delta w_7)), \quad (31)$$

where $5 \leq i \leq 6$.

Case 5 ($k_7 = 7$):

$$C(\mu) \leq C(\mu - (w_7 + 7(w_7 - w_6))) + C(\mu - (8w_7 + 26\Delta w_7)). \quad (32)$$

We have derived recurrences for all branching operations in algorithm $\text{mis7}(G)$ except for Step 5 which invokes algorithms $\text{mis6}(G)$. To determine the largest branching factor to algorithm $\text{mis7}(G)$ analogously with the previous section, we combine all the above recurrences with the weight setting used for $\text{mis6}(G)$. By Lemma 4, we include the following four constraints into the current set of recurrences.

$$\begin{aligned} C(\mu) &\leq 1.18922 \frac{w_3^{(6)}}{w_3} \mu, & C(\mu) &\leq 1.18922 \frac{w_4^{(6)}}{w_4} \mu, \\ C(\mu) &\leq 1.18922 \frac{w_5^{(6)}}{w_5} \mu, & \text{and } C(\mu) &\leq 1.18922 \frac{w_6^{(6)}}{w_6} \mu, \end{aligned} \quad (33)$$

where $w_3^{(6)} = 0.49969$, $w_4^{(6)} = 0.76163$, $w_5^{(6)} = 0.92401$ and $w_6^{(6)} = 1$.

The above assumptions and recurrences together with (33) generate the constraints in our quasiconvex program. By solving the quasiconvex program, we get an upper bound 1.19698 on the branching factor for all recurrences by setting vertex weights as

$$w_i = \begin{cases} 0 & \text{for } i = 0, 1 \text{ and } 2 \\ 0.65077 & \text{for } i = 3 \\ 0.78229 & \text{for } i = 4 \\ 0.89060 & \text{for } i = 5 \\ 0.96384 & \text{for } i = 6 \\ 1 & \text{for } i = 7 \\ w_7 + (i - 7)(w_7 - w_6) & \text{for } i \geq 8. \end{cases} \quad (34)$$

This verifies Lemma 10 with $\theta = 7$.

9 Analysis of $\text{mis8}(G)$

Recall that for MIS-8, we assume $w_0 = w_1 = w_2 = 0 \leq w_3 \leq w_4 \leq w_5 \leq w_6 \leq w_7 \leq w_8 = 1 \leq w_9 \leq \dots$, and the values of w_3, w_4, w_5, w_6 and w_7 will be determined after we analyze how the measure changes after each step of the algorithm.

To simplify analysis, we assume that

$$26\Delta w_8 \leq w_3. \quad (35)$$

To ensure that contracting vertices never increase the measure, we impose the next constraint.

$$w_i + w_j \geq w_{i+j-2}, \quad 3 \leq i, j \leq 8. \quad (36)$$

Next we analyze each step of the algorithm.

Step 1: We can see that Lemma 13 still holds in $\text{mis8}(G)$ and then the measure of the graph will not increase after Step 1 of $\text{mis8}(G)$.

Step 2: Using recurrences (2) for branching on a vertex of degree ≥ 9 in Step 2, we get recurrences:

$$\begin{aligned} C(\mu) &\leq C(\mu - (w_d + \sum_{i=3}^d k_i \Delta w_i)) + C(\mu - (w_d + \sum_{i=3}^d k_i w_i + \Delta(\overline{N[v]}))) \\ &\leq \max_{3 \leq i \leq 9} [C(\mu - (w_9 + 9\Delta w_i)) + C(\mu - (w_9 + 9w_i + 16\Delta w_8))]. \end{aligned} \quad (37)$$

Step 3: We consider branching on an optimal short edge vv' in Step 3. We see that $|N(v) \cap N(v')| \geq 7$ cannot occur, otherwise v would dominate v' .

(i) For the case of $|N(v) \cap N(v')| = 6$, we get the following recurrences for the branch (analogously with Case (i) in Section 7.3)

$$\begin{aligned} C(\mu) &\leq C(\mu - (2w_8 + 6(w_i - w_{i-2}) + 2\Delta w_8)) \\ &\quad + C(\mu - (2w_8 + 6w_i + 2\Delta w_8)) \quad (i = 3, 4, 5, 6, 7). \end{aligned} \quad (38)$$

(ii) For the case of $|N(v) \cap N(v')| = 5$, we get the following recurrences

$$\begin{aligned} C(\mu) &\leq C(\mu - (2w_8 + 5(w_i - w_{i-2}) + 4\Delta w_j)) \\ &\quad + C(\mu - (2w_8 + 5w_i + (5 + p)\Delta w_8 - 4(w_{j+1} - w_j))), \end{aligned} \quad (39)$$

where $4 \leq i \leq 8$ ($p = 5$ for $i = 8$ and $p = 0$ for $i = 4, 5, 6$ and 7) and $3 \leq j \leq 8$; and

$$C(\mu) \leq C(\mu - (2w_8 + 10w_3 + 4\Delta w_j)) + C(\mu - (2w_8 + 5w_3 + 4\Delta w_7 - 4(w_{j+1} - w_j))), \quad (40)$$

where $3 \leq j \leq 8$.

(iii) For the case of $|N(v) \cap N(v')| = 4$, we get the following recurrences

$$C(\mu) \leq C(\mu - (2w_8 + 4(w_i - w_{i-2}) + 6\Delta w_j)) + C(\mu - (2w_8 + 4w_i + (4+p)\Delta w_8 - 6(w_{j+1} - w_j))), \quad (41)$$

where $4 \leq i \leq 8$ ($p = 8$ for $i = 8$; $p = 4$ for $i = 7$; and $p = 0$ for $i = 4, 5$ and 6) and $3 \leq j \leq 8$; and

$$C(\mu) \leq C(\mu - (2w_8 + 8w_3 + 6\Delta w_j)) + C(\mu - (2w_8 + 4w_3 + 4\Delta w_7 - 6(w_{j+1} - w_j))), \quad (42)$$

where $3 \leq j \leq 8$.

Step 4: Using recurrences (2) for branching on an optimal vertex of degree 8 in Step 4, we can get recurrences:

$$C(\mu) \leq C(\mu - (w_8 + \sum_{i=3}^8 k_i \Delta w_i)) + C(\mu - (w_8 + \sum_{i=3}^8 k_i w_i + \lambda_8(k_8))), \quad (43)$$

for all nonnegative integers $(k_3, k_4, k_5, k_6, k_7, k_8)$ with $k_3 + k_4 + k_5 + k_6 + k_7 + k_8 = 8$ and

$$\lambda_8(k_8) = \begin{cases} (16 + 2k_8)\Delta w_8 & \text{if } k_8 \leq 7 \\ 36\Delta w_8 & \text{if } k_8 = 8. \end{cases} \quad (44)$$

The correctness of the above recurrences relies on the following lemma, which corresponds to Lemma 17.

Lemma 18 *Let v be an optimal degree-8 vertex in Step 4 of $\text{mis8}(G)$. Then $\Delta(\overline{N[v]}) \geq \lambda_8(k_8)$.*

Proof. If $N^*(v) = \emptyset$. Then each neighbor of v has at least two neighbors in $N_2(v)$ and each degree-8 neighbor of v has at least four neighbors in $N_2(v)$. We obtain $\Delta(\overline{N[v]}) \geq f_v \Delta w_8 \geq (16 + 2k_8)\Delta w_8$. By the definition of optimal vertices, we know that when $k_8 = 8$, it holds $f_v \geq 36$ and then $\Delta(\overline{N[v]}) \geq 36\Delta w_8$.

We see that the statement of Lemma 15(ii) still holds for $\theta = 8$ even after replacing ‘ Δw_6 ’ with ‘ Δw_8 ’ in it. This implies that if $N^*(v) \neq \emptyset$, then $\Delta(\overline{N[v]}) \geq \min\{2w_3, w_3 + 2(\delta(v) - 3)\Delta w_8\} \geq w_3 + 10\Delta w_8$, which is larger than any of $(16 + 2k_8)\Delta w_8$ and $36\Delta w_8$ by (35). \blacksquare

Step 5: In Step 5, the algorithm invokes $\text{mis7}(G)$. Analogously with the previous section, we include the following five constraints into the current set of recurrences.

$$C(\mu) \leq 1.19698 \frac{w_3^{(7)}}{w_3} \mu, C(\mu) \leq 1.19698 \frac{w_4^{(7)}}{w_4} \mu, C(\mu) \leq 1.19698 \frac{w_5^{(7)}}{w_5} \mu, \quad (45)$$

$$C(\mu) \leq 1.19698 \frac{w_6^{(7)}}{w_6} \mu, \text{ and } C(\mu) \leq 1.19698 \frac{w_7^{(7)}}{w_7} \mu,$$

where $w_3^{(7)} = 0.65077$, $w_4^{(7)} = 0.78229$, $w_5^{(7)} = 0.89060$, $w_6^{(7)} = 0.96384$ and $w_7^{(7)} = 1$.

After solving the quasiconvex program, we get an upper bound 1.19951 on the branching factor for all recurrences by setting vertex weight:

$$w_i = \begin{cases} 0 & \text{for } i = 0, 1 \text{ and } 2 \\ 0.65844 & \text{for } i = 3 \\ 0.78844 & \text{for } i = 4 \\ 0.88027 & \text{for } i = 5 \\ 0.95345 & \text{for } i = 6 \\ 0.98839 & \text{for } i = 7 \\ 1 & \text{for } i = 8 \\ w_8 + (i - 8)(w_8 - w_7) & \text{for } i \geq 9. \end{cases} \quad (46)$$

This verifies Lemma 10 with $\theta = 8$.

10 Proof of Lemma 9

We prove Lemma 9 by revealing some structural properties of graphs of maximum degree 6, 7 and 8. Recall that, for a vertex v , f_v denotes the number of edges between $N(v)$ and $N_2(v)$, and e_v denotes the number of edges in the graph $G[N(v)]$. For each neighbor $u \in N(v)$, the outer-degree (resp., inner-degree) of u at v is $|N(u) \cap N_2(v)|$ (resp., $|N(u) \cap N(v)|$).

10.1 Graphs of maximum degree 6

The existence of optimal vertices in a reduced graph G of maximum degree 6 without short edges follows from Lemma 19. When there is not short edge in a reduced graph with maximum degree 6, we see that for each degree-6 vertex v in G , the inner-degree of any vertex in $N(v)$ at v is at most 2. Such a graph can have the following types of vertices.

Lemma 19 *Let G be a graph of maximum degree 6 and minimum degree ≥ 3 such that for every degree-6 vertex v , the inner-degree of each neighbor $u \in N(v)$ at v is at most 2. If G is not the line graph of a 4-regular graph, then there is a degree-6 vertex v that satisfies one of the following:*

- $k_3 \geq 1$ or $k_6 \leq 3$;
- $k_6 = 4$ and $k_5 \leq 1$;
- $k_6 = 4, k_5 = 2$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 17$;
- $k_6 = 5, k_4 = 1$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 18$;
- $k_6 = 5, k_5 = 1$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 19$; and
- $k_6 = 6$ and $f_v + (f_v - |N_2(v)|) + q_v \geq 22$.

Proof. Observe that f_v is the sum of the out-degree of neighbors of v and

$$e_v \leq 6$$

holds, since the inner-degree of each neighbor at v is at most 2. We assume that G has no vertex that satisfies one of “ $k_3 \geq 1$ or $k_6 \leq 3$ ” and “ $k_6 = 4$ and $k_5 \leq 1$.” We consider several cases.

Case 1. There is a degree-6 vertex v with $k_6 = 5$ and $k_4 = 1$: Assume that $f_v + (f_v - |N_2(v)|) + q_v \leq 16$. Since $f_v \geq 16$ by $e_v \leq 6$, we see that $f_v - |N_2(v)| = q_v = 0$ and the degree-4 neighbor u of v is adjacent to a degree-6 vertex $z \in N(u) \cap N_2(v)$. Now $N(z) - \{u\}$ contains only degree-6 vertices, since z has already one degree-4 neighbor. Note that z is not adjacent to any vertex in $N[v] - \{u\}$, and then the outer-degree of vertex u at z is three. This implies that $f_z \geq 3 \times 5 + 3 = 18$ and v is a vertex satisfying the condition in the lemma.

In what follows, we further assume that there is no degree-6 vertex with $k_6 = 5$ and $k_4 = 1$. We choose a degree-6 vertex v with minimum e_v such that $k_6 < 6$ (if possible) and then the maximum component in $G[N(v)]$ is maximized.

Case 2. $e_v \leq 4$: In this case, we are done because $f_v \geq 16 + 2 \times (6 - e_v) = 20$ for $k_6 = 4$ and $k_5 = 2$; $f_v \geq 17 + 4 = 21$ for $k_6 = 5$ and $k_5 = 1$; and $f_v \geq 18 + 4 = 22$ for $k_6 = 6$.

Case 3. $e_v = 5$: In this case, $f_v \geq 16 + 2 \times (6 - e_v) = 18$ for $k_6 = 4$ and $k_5 = 2$; $f_v \geq 17 + 2 = 19$ for $k_6 = 5$ and $k_5 = 1$; and $f_v \geq 18 + 2 = 20$ for $k_6 = 6$. We only need to consider the case of $k_6 = 6$, and assume that $f_v + (f_v - |N_2(v)|) + q_v \leq 21$, where we have $f_v - |N_2(v)| + q_v \leq 1$ by $f_v \geq 20$. Observe that $G[N(v)]$ is either a path of length 5 or a disjoint union of a path of i and a cycle of length $5 - i$ ($i = 0, 1, 2$). We distinguish two subcases.

(i) $G[N(v)]$ contains a path of length four $u_1 u_2 u_3 u_4 u_5$ (possibly a cycle of length 4 with $u_1 = u_5$): We show that u_3 satisfies the lemma. By $f_v - |N_2(v)| + q_v \leq 1$, at most one of u_2 and u_4 can be adjacent to a vertex in $N(u_3) \cap N_2(v)$; i.e., the inner-degree of u_i ($i = 2, 4$) at u_3 is at most 1, implying that $e_{u_3} \leq 5$ and u_3 has only degree-6 neighbors by our choice of v . Note that

$f_{u_3} - |N_2(u_3)| \geq 2$, since u_1 and u_5 are common neighbors of two neighbors in $N(u_3)$. This means that $f_{u_3} + (f_{u_3} - |N_2(u_3)|) \geq 20 + 2 = 22$.

(ii) $G[N(v)]$ consists of a path $u_1u_2u_3$ and a triangle: By $f_v - |N_2(v)| \leq 1$, we see that for $i = 1$ or 3 (say $i = 1$), there is no edge between $N(u_i) \cap N_2(v)$ and $N(v) - \{u_i\}$. In this case, the inner-degree of each of v and u_2 at u_1 is 1, implying that $e_{u_1} \leq 5$ (hence $e_{u_1} = 5$). Hence u_1 has only degree-6 neighbors by our choice of v . However, in this case, $G[N(u_1)]$ must be a union of single edge vu_2 and a cycle of length 4. Then u_1 satisfies the condition of (i).

Case 4. $e_v = 6$. In this case, $f_v \geq 16$ for $k_6 = 4$ and $k_5 = 2$; $f_v \geq 17$ for $k_6 = 5$ and $k_5 = 1$; and $f_v \geq 18$ for $k_6 = 6$.

We first consider the case of $k_6 = 4$ and $k_5 = 2$. Assume $f_v + (f_v - |N_2(v)|) + q_v \leq 16$ (otherwise we are done), which implies that $f_v - |N_2(v)| = q_v = 0$. Let $u \in N(v)$ be a degree-5 neighbor of v , and $z \in N(u) \cap N_2(v)$ be a neighbor of u not in $N[v]$, where $\delta(z) = 6$ (by $q_v = 0$) and z is not adjacent to any other vertices in $N(v) - \{u\}$ (by $f_v - |N_2(v)| = 0$). Hence the inner-degree of u at z is at most 1, and $e_z \leq 5$, contradicting our choice of vertex v .

Assume that v satisfies “ $k_6 = 5$ and $k_5 = 1$ ” or “ $k_6 = 6$.” Now $G[N(v)]$ with $e_v = 6$ is either a cycle of length 6 or a disjoint union of two triangles. We distinguish two subcases.

(i) $G[N(v)]$ is a cycle of length six $u_1u_2u_3u_4u_5u_6$, where u_1 is assumed to be of degree 5 if $k_5 = 1$: By our choice of v , it also holds $e_{u_2} = 6$, implying that each of u_1 and u_3 is adjacent to a vertex in $N(u_2) \cap N_2(v)$. Analogously each of u_4 and u_6 is adjacent to a vertex in $N(u_5) \cap N_2(v)$. Hence we have $f_v - |N_2(v)| \geq 4$ and $f_v + (f_v - |N_2(v)|)$ is at least $16 + 4 = 20$ for “ $k_6 = 5$ and $k_5 = 1$ ” and $18 + 4 = 22$ for $k_6 = 6$, as required.

(ii) $G[N(v)]$ is a disjoint union of two triangles: In this case, v satisfies “ $k_6 = 5$ and $k_5 = 1$ ” only, because otherwise by our choice G would be a 6-regular graph such that for each vertex v' in it, $G[N(v)]$ is a disjoint union of two triangles and then G would be the line graph of a 4-regular graph. Assume that $f_v + (f_v - |N_2(v)|) + q_v \leq 18$ (otherwise we are done), where $(f_v - |N_2(v)|) + q_v \leq 1$. Let $u \in N(v)$ be the degree-5 neighbor of v , and let $\{z_1, z_2\} = N(u) \cap N_2(v)$. By $q_v \leq 1$, one of z_1 and z_2 (say z_1) is of degree 6. Since $e_{z_1} = 6$ by our choice of v , the inner-degree of u at z_1 is 2, and hence z_1 is adjacent to a neighbor in $N(v) \cap N(u)$. Now $f_v - |N_2(v)| = 1$ and hence $q_v = 0$. The vertex z_2 also needs to be adjacent to a neighbor in $N(v) \cap N(u)$, indicating $f_v - |N_2(v)| \geq 2$, a contradiction. \blacksquare

10.2 Graphs of maximum degree 7

To prove the existence of optimal vertices in graphs of maximum degree 7, we investigate the structure of 7-regular graphs.

A vertex v is called a (j, k) -clique-type vertex if the induced graph $G[N(v)]$ is a disjoint union of a j -clique and a k -clique, where $j + k = \delta(v)$.

Lemma 20 *Let G be a 7-regular graph such that the inner-degree of each neighbor $u \in N(v)$ of every vertex v is at most 3. If G is not the line graph of a $(4, 5)$ -bipartite graph, then it has a vertex v such that $f_v + (f_v - |N_2(v)|) \geq 26$.*

Proof. Now $f_v = 42 - 2e_v$ is an even number. Assume that for every vertex v , $f_v + (f_v - |N_2(v)|) \leq 25$ otherwise we are done. Before we prove that G is the line graph of a $(4, 5)$ -bipartite graph, we first show four properties (P0), (P1), (P2) and (P3) on a vertex v in G .

(P0) *It holds that $f_v \in \{22, 24\}$ and $e_v \geq 9$.*

Now the inner-degree of each neighbor $u \in N(v)$ at v is at most 3, i.e., the outer-degree of u of v is at least 3 in a 7-regular graph, which implies $f_v \geq 3 \times 7 = 21$ and hence $f_v \geq 22$ by parity. By $f_v = 42 - 2e_v$, we see that $f_v + (f_v - |N_2(v)|) \leq 25$ implies $f_v \leq 24$ and $e_v \geq 9$.

(P1) *If a vertex v is not $(3, 4)$ -clique-type, then there is no 4-clique in $G[N(v)]$.*

If $G[X]$ is a 4-clique for some subset $X \subseteq N(v)$, then the inner-degree of a neighbor $u \in X$ is already 3 and the remaining $e_v - 6 \geq 9 - 6 = 3$ edges must form a triangle in $N(v) - X$, indicating that v would be (3,4)-clique-type.

(P2) *If there is no 4-clique in $G[N(v)]$, then there are four neighbors $u_1, u_2, u_3, u_4 \in N(v)$ such that there is at least one edge between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$.*

Since $f_v \leq 24$ by (P0), there are at least four neighbors $u_1, u_2, u_3, u_4 \in N(v)$, each of which has outer-degree 3 at v (i.e., $|N(u_i) - N_2(v)| = 4$). If there is no edge between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$, then $e_{u_i} \geq 9$ by (P0) implies that $N(u_i) \cap N(v)$ and $N(u_i) - N_2(v)$ induce a 3-clique and 4-clique, respectively (where $N(u_i) \cap N(v)$ induces a 3-clique), which contradicts that $\{u_i\} \cup (N(u_i) \cap N(v))$ does not induce a 4-clique.

(P3) *If there is exactly one edge between X and $X - N(v)$ for some set X of four vertices in $N(v)$, then v is not (3,4)-clique-type and $f_v \geq 24$.*

If $f_v \leq 22$ (i.e., $e_v \geq 10$) then $G[X]$ needs to be a 4-clique and the inner-degree of some vertex in X at v would be 4.

We are ready to prove the lemma by using (P0), (P1), (P2) and (P3). If the graph is not the line graph of a (4,5)-bipartite graph, then we can always choose a vertex v that is not (3,4)-clique-type so that $f_v \in \{22, 24\}$ is maximized. By (P1) and (P2), there are four neighbors $u_i \in N(v)$, $i = 1, 2, 3, 4$ such that there is at least one edge between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$.

(i) For each i , there are two such edges: Then we get $f_v - |N_2(v)| \geq 4$ and it would hold $f_v + (f_v - |N_2(v)|) \geq 22 + 4 = 26$, a contradiction.

(ii) For some i , there is exactly one such edge: In this case, we get $f_v - |N_2(v)| \geq 2$. Since u_i satisfies the condition of (P3), it holds $f_{u_i} \geq 24$, and by the choice of v , we have $f_v \geq f_{u_i} \geq 24$. Hence it would hold $f_v + (f_v - |N_2(v)|) \geq 24 + 2 = 26$, a contradiction. \blacksquare

Lemma 21 *Let G be a reduced graph of maximum degree 7. Assume that G has no short edges. Then G has at least one optimal vertex.*

Proof. We assume that every degree-7 vertex v has at least six neighbors of degree 7 and satisfies $N^*(v) = \emptyset$, otherwise the lemma holds. Each neighbor of v is adjacent to at least two vertices in $N_2(v)$ (since $N^*(v) = \emptyset$) and each degree-7 neighbor of v is adjacent to at least three vertices in $N_2(v)$ (since here is no short edge). Hence $f_v \geq 20$. If v is adjacent to a degree-3 vertex, then v is an optimal vertex by the definition of optimal vertices. If the graph is a 7-regular graph, then there is an optimal vertex by Lemma 20. Otherwise, we can always find a vertex v with $k_7 = 6$ and $k_3 = 0$.

Let u_1 be the neighbor of v such that $4 \leq \delta(u_1) \leq 6$. If u_1 is not adjacent to any other vertex in $N(v)$, then $f_v \geq \delta(u_1) - 1 + 3 \times 6 = \delta(u_1) + 17$ and v would be an optimal vertex. Otherwise, u_1 is adjacent to a degree-7 vertex $u_2 \in N(v)$. If u_2 has outer-degree at least 5 at v , then $f_v \geq 5 + 3 \times 5 + 2 = 22$ and v would be an optimal vertex. We can assume that u_2 has outer-degree 3 or 4 at v . If there are at least two edges between $N(u_2) \cap N(v)$ and $N(u_2) \cap N_2(v)$, then $f_v - |N_2(v)| \geq 2$ and v would be an optimal vertex. Otherwise, there is at most one edge between $N(u_2) \cap N(v)$ and $N(u_2) \cap N_2(v)$. Since $\{|N(u_2) \cap N(v)|, |N(u_2) \cap N_2(v)|\} = \{3, 4\}$, then $e_{u_2} \leq 10$ and $f_{u_2} \geq 22$. Note that u_2 is also adjacent to a vertex u_1 with degree < 7 . Then u_2 will be an optimal vertex for this case. \blacksquare

10.3 Graphs of maximum degree 8

To prove the existence of optimal vertices in graphs of maximum degree 8, we investigate the structure of 8-regular graphs.

First of all, we consider 8-regular graph such that the inner-degree of each neighbor $u \in N(v)$ of every vertex v is at most 3. Now $f_v = 56 - 2e_v$ is an even number. This properties will be

used in the following lemmas several times. A vertex v is called *bridge-type* if $G[N(v)]$ contains a bridge u_1u_2 between $X \subseteq N(v)$ and $N(v) - X$ such that (i) $|X| = |N(v) - X|$; or (ii) the inner-degree of each u_i at v is 3.

Lemma 22 *Let G be a 8-regular graph such that the inner-degree of each neighbor $u \in N(v)$ of every vertex v is at most 3. If G contains a bridge-type vertex, then there is a vertex v such that $f_v + (f_v - |N_2(v)|) \geq 36$.*

Proof. Let v a bridge-type vertex v , and u_1u_2 be the bridge of v between X_1 and $X_2 = N(v) - X$, where $u_i \in X_i$ and $|X_1| \leq |X_2|$ without loss of generality.

We first prove that $f_v \geq 34$. Note that $|X_1| \geq 3$, since otherwise the inner-degree of u_1 at v would be at most $|X_1| - 1 \leq 2$. When $|X_1| = 3$, the inner-degree of each of the two vertices in $X_1 - \{u_1\}$ is at most 2 at v , since u_1u_2 is a bridge in $G[N(v)]$. When $|X_1| = |X_2| = 4$, $G[X_i]$ contains at least one pair of non-adjacent vertices, since otherwise the inner-degree of u_i would be four, and X_i contains at least one vertex whose inner-degree at v is at most 2. In any case of $|X_1| \in \{3, 4\}$, $N(v)$ contains at least two neighbors whose outer-degree at v is at least 5. This implies that $f_v \geq 6 \times 4 + 2 \times 5 = 34$.

We further assume that $N(v)$ consists of two (resp., six) neighbors whose inner-degree at v are 2 (resp., 3), since otherwise $f_v \geq 35$ (hence $f_v \geq 36$ by parity) and we are done. Note that there are at least three neighbors in $N(v)$ whose inner-degree are 2 if $|X_1| = 3$ (2 in X_1 and 1 in X_2). So it holds $|X_1| = 4$ and the inner-degree of u_i at v is 3.

The outer-degree of u_i ($i = 1, 2$) at v is 4 (i.e., $|N(u_i) \cap N[v]| = 4$), and the induced graph $G[N(u_i) \cap N[v]]$ contains at least two pairs of non-adjacent vertices (since there is only one edge between $N[u_i] \cap N(v)$ and $N(v) - N[u_i] \cap N(v)$). If there is no edge between $N(u_i) \cap N[v]$ and $N(u_i) \cap N_2(v)$, then $e_{u_i} \leq 10$ and $f_{u_i} \geq 56 - 2 \times 10 = 36$. Hence we can assume that u_i has a common neighbor in $N_2(v)$ with a vertex in $N(v) - \{u_i\}$.

Let u'_i ($i = 1, 2$) be the two neighbors in $N(v)$ whose inner-degree at v is 2. If there is no edge between $N(u'_i) \cap N[v]$ and $N(u_i) \cap N_2(v)$, then each vertex in $N(u'_i) \cap N[v]$ has outer-degree 5 at u'_i . We get that $f_{u'_i} \geq 3 \times 5 + 5 \times 4 = 35$ (hence $f_{u'_i} \geq 36$ by parity). Hence we can assume that u'_i has a common neighbor in $N_2(v)$ with a vertex in $N(v) - \{u'_i\}$.

Thus there are four vertices $\{u_1, u_2, u'_1, u'_2\}$ each of who has a common neighbor in $N_2(v)$ with another vertex in $N(v)$, which implies that $f_v - |N_2(v)| \geq 4/2 = 2$, and we have that $f_v + (f_v - |N_2(v)|) \geq 34 + 2 = 36$. \blacksquare

Recall that a vertex v is (j, k) -clique-type if the induced graph $G[N(v)]$ is a disjoint union of a j -clique and a k -clique, where $j+k = \delta(v)$. A vertex v is called *semi-clique-type* if a set $X \subseteq N(v)$ induces a 4-clique, there is no edge between X and $N(v) - X$, and $X \subseteq N(v)$ contain at least one pair of non-adjacent vertices. Hence a semi-clique-type vertex is not $(4, 4)$ -clique-type.

Lemma 23 *Let G be a 8-regular graph such that the inner-degree of each neighbor $u \in N(v)$ of every vertex v is at most 3. If G contains a semi-clique-type vertex, then there is a vertex v such that $f_v + (f_v - |N_2(v)|) \geq 36$.*

Proof. Let v be a semi-clique-type vertex, and assume that $X \subseteq N(v)$ induce a 4-clique. Clearly $f_v \geq 56 - 2 \times 11 = 34$ (since $e_v \leq 11$). Assume that $N(v) - X$ contains only one pair of non-adjacent vertices u_1 and u_2 , since otherwise we obtain $f_v \geq 56 - 2 \times 10 = 36$. Let $N(v) - X = \{u_1, u_2, u_3, u_4\}$. If each of u_3 and u_4 is adjacent to a vertex in $N(u_1) - N[v]$, then we have $f_v - |N_2(v)| \geq 2$, and $f_v + (f_v - |N_2(v)|) \geq 34 + 2 = 36$. If none of u_3 and u_4 is adjacent to any vertex in $N(u_1) - N[v]$, then all of v , u_3 and u_4 have inner-degree 2 at u_1 , and this implies that $f_{u_1} \geq 36$. Finally if exactly one of u_3 and u_4 is adjacent to a vertex in $N(u_1) - N[v]$, then u_1 is bridge-type, and $f_{u_1} + (f_{u_1} - |N_2(u_1)|) \geq 36$ by Lemma 22. \blacksquare

Lemma 24 *Let G be a 8-regular graph such that the inner-degree of each neighbor $u \in N(v)$ of every vertex v is at most 3. Assume that G is not the line graph of a 5-regular graph. Then G has a vertex v such that $f_v + (f_v - |N_2(v)|) \geq 36$.*

Proof. By Lemma 22 and Lemma 23, it suffices to show that there is a desired vertex or a bridge-type or semi-clique-type vertex. Let v be a vertex that is not (4,4)-clique-type. This vertex always exists since the graph is not the line graph of a 5-regular graph. Assume that v is not bridge-type or semi-clique-type. Hence no four neighbors of v induce a 4-clique.

Assume $f_v \leq 34$, otherwise we are done. This implies that there are at least six neighbors $u_i \in N(v)$, $i = 1, \dots, 6$ each of which has outer-degree 4 at v (i.e., $|N[u_i] \cap N(v)| = 4$).

If for each i , there are at least two edges between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$, then we have $f_v - |N_2(v)| \geq 12/2 = 6$, and $f_v + (f_v - |N_2(v)|) \geq 32 + 6 > 36$. Assume that for some i , there is at most one edge between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$. If there is exactly one edge between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$, then u_i is bridge-type. Assume that there is no edge between $N(u_i) \cap N(v)$ and $N(u_i) \cap N_2(v)$. Recall that no four vertices in $N(v)$ induce a 4-clique. Hence $N(u_i) \cap N[v]$ is not a 4-clique either, and this means that u_i is semi-clique-type. ■

Lemma 25 *Let G be a reduced graph of maximum degree 8. If G has no short edges, then G has at least one optimal vertex.*

Proof. This lemma follows from the definition of optimal vertices and Lemma 24 directly. If the graph G is not a 8-regular graph, we can always find a degree-8 vertex such that $k_8 < 8$, which is an optimal vertex by the definition. Otherwise, G is a 8-regular graph. Since G has no short edges, the inner-degree of each neighbor $u \in N(v)$ of every vertex v is at most 3. Then by Lemma 24 there is either an optimal vertex or the graph is the line graph of a 5-regular graph. However, the later case is impossible, since the line graph of a 5-regular graph must have been reduced by the reduction rules. ■

11 Concluding Remarks

Before the measure-and-conquer method was developed, most fast algorithms for the maximum independent set problem consisted of a large number of branching rules, which may make the algorithms impractical and hard to analyze. The measure-and-conquer method allows us to design simple algorithms for the maximum independent set problem probably with an aid of sophisticated analysis. With this method, we get the recurrence (2) for branching on a vertex v of maximum degree d , which usually becomes the worst case of algorithms to any MIS- θ ($3 \leq \theta \leq 8$). To analyze (2), we need to do both of (i) checking all possible neighbor-degree (k_3, k_4, \dots, k_d) of the neighbors of v ; and (ii) deriving lower bounds on the term $\Delta(\overline{N[v]})$.

For (i), the previous papers either to try to reduce the number of cases to be checked by a relaxed argument (and then get worse recurrences) or list up a huge number of recurrences for all possible combinations (which may not be easy to check by hand). In this paper, we devised a new lemma (Lemma 6) that can reduce the number of cases to a quite small number without losing the optimality of branching factors. In the branch-and-reduce paradigm, this is useful to simplify analysis of algorithms and can make a design process of fast algorithms much easier.

For (ii), there are many techniques used to derive good bounds on $\Delta(\overline{N[v]})$. With the reduction rule by domination, Fomin *et al.* [4] got that $\Delta(\overline{N[v]}) \geq d\Delta w_d$. With branching on vertices with satellites (which is extended to unconfined vertices later in [16]), Kneis *et al.* [8] showed $\Delta(\overline{N[v]}) \geq 2d\Delta w_d$ in the worst case $k_d = d$ of (2) (this is also used in Bourgeois *et al.*'s algorithm [1]). In this paper, by using the new branching rule to short edges, for the worst case of (2), we improved the bounds on $\Delta(\overline{N[v]})$ to $3d\Delta w_d$ for $d = 6, 7$ and to $4d\Delta w_d$ for $d \geq 8$,

respectively. By choosing an optimal vertex whose existence is ensured by a graph theoretical argument, we further increased the bound on $\Delta(N[v])$ to $(4d + 4)\Delta w_d$ for the case of “ $d = 8$ and $k_d = d$,” which is the final worst case in our algorithms. Branching on a degree-8 vertex v with eight degree-8 neighbors and 36 edges between $N(v)$ and $N_2(v)$ (i.e., $f_v = 36$ for $k_8 = 8$) is one of the crucial bottlenecks in our algorithm for MIS now.

Acknowledgements

A preliminary version of this paper with a result of $1.2002^n n^{O(1)}$ was presented in the 24th international symposium on algorithms and computation (ISAAC 2013) and appeared as [19]. In the full version, we further reduce several bottlenecks by refining the definition of “optimal vertices” and then finally break the barrier of 1.2 in the base of the running time. Part of the work was done when the first author visited Kyoto University in 2013.

References

- [1] Bourgeois, N., Escoffier, B., Paschos, V. T., van Rooij, J. M. M., Fast algorithms for max independent set, *Algorithmica* 62(1-2), (2012) 382–415.
- [2] Chen, J., Kanj, I.A., Xia, G.: Improved upper bounds for vertex cover. *Theoretical Computer Science* 411(40-42) (2010) 3736–3756
- [3] Eppstein D.: Quasiconvex analysis of multivariate recurrence equations for backtracking algorithms. *ACM Transactions on Algorithms* 2(4) (2006) 492–509
- [4] Fomin, F. V., Grandoni, F., Kratsch, D.: A measure & conquer approach for the analysis of exact algorithms. *J. ACM* 56(5) (2009) 1–32
- [5] Fomin, F. V., Kratsch, D.: *Exact Exponential Algorithms*, Springer (2010)
- [6] Fürer, M.: A faster algorithm for finding maximum independent sets in sparse graphs. In: *LATIN 2006*. LNCS 3887, (2006) 491–501
- [7] Jian, T.: An $O(2^{0.304n})$ algorithm for solving maximum independent set problem. *IEEE Transactions on Computers* 35(9) (1986) 847–851
- [8] Kneis, J., Langer, A., Rossmanith, P.: A fine-grained analysis of a simple independent set algorithm. In *FSTTCS 2009*. V. 4 LIPIcs., (2009) 287–298
- [9] Razgon, I.: Faster computation of maximum independent set and parameterized vertex cover for graphs with degree 3. *J. of Disc. Alg.* 7(2) (2009) 191–212
- [10] Robson, J.: Algorithms for maximum independent sets. *J. of Algorithms* 7(3) (1986) 425–440
- [11] Robson, J.: Finding a maximum independent set in time $O(2^{n/4})$. Technical Report 1251-01, LaBRI, Univsersite Bordeaux I (2001)
- [12] Tarjan, R., Trojanowski, A.: Finding a maximum independent set. *SIAM J. on Computing* 6(3) (1977) 537–546
- [13] West, D.: *Introduction to Graph Theory*. Prentice Hall. 1996

- [14] Woeginger, G.J.: Exact algorithms for NP-hard problems: a survey. In: Combinatorial Optimization–Eureka! You shrink! LNCS 2570, (2003) 185–207
- [15] Xiao, M.: A simple and fast algorithm for maximum independent set in 3-degree graphs. In: M. Rahman and S. Fujita: WALCOM 2010, LNCS 5942, (2010) 281–292
- [16] Xiao, M., Nagamochi, H.: Confining sets and avoiding bottleneck cases: A simple maximum independent set algorithm in degree-3 graphs. Theoretical Computer Science **469** (2013) 92–104
- [17] Xiao, M., Nagamochi, H.: A refined algorithm for maximum independent set in degree-4 graphs. Technical report 2013-002, Kyoto University (2013) URL: <http://www.amp.i.kyoto-u.ac.jp/tecrep/abst/2013/2013-002.html>
- [18] Xiao, M., Nagamochi, H.: An exact algorithm for maximum independent set in degree-5 graphs. In: FAW-AAIM 2013, LNCS 7924, (2013) 72–83
- [19] Xiao, M. and Nagamochi, H.: Exact algorithms for maximum independent set. In: ISAAC 2013. LNCS 8283 (2013) 328–338
- [20] URL: <http://www.tcs.rwth-aachen.de/independentset/>