

オペレーションズ・リサーチ

3

$\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{b} \in \mathbb{R}^m$, $\mathbf{C} \in \mathbb{R}^{n \times n}$ とする. パラメータ $\mathbf{x} = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$ をもつ次の非線形計画問題を考える.

$$\begin{aligned} \text{P}(\mathbf{x}): \quad & \text{Minimize} \quad \sum_{i=1}^n (\mathbf{z}^i)^\top \mathbf{z}^i + \mathbf{y}^\top \mathbf{y} + \mathbf{x}^\top \mathbf{C} \mathbf{x} \\ & \text{subject to} \quad \mathbf{y} - \sum_{i=1}^n x_i \mathbf{z}^i = \mathbf{A} \mathbf{x} - \mathbf{b} \end{aligned}$$

ここで, $\text{P}(\mathbf{x})$ の決定変数は $\mathbf{y}, \mathbf{z}^i \in \mathbb{R}^m$ ($i = 1, \dots, n$) である. また, $^\top$ は転置記号を表す. さらに, 任意の \mathbf{x} に対して, 問題 $\text{P}(\mathbf{x})$ の最適値が定義されているとし, その最適値を $f(\mathbf{x})$ と表す.

以下の問いに答えよ.

- (i) 問題 $\text{P}(\mathbf{x})$ のカルーシュ・キューン・タッカー条件 (Karush-Kuhn-Tucker 条件) を書け.
- (ii) 問題 $\text{P}(\mathbf{x})$ の目的関数が, $\mathbf{y}, \mathbf{z}^i \in \mathbb{R}^m$ ($i = 1, \dots, n$) に対して凸であることを示せ.
- (iii) \mathbf{C} を正定値対称行列と仮定し, 次の最適化問題を考える.

$$\begin{aligned} \text{P1:} \quad & \text{Minimize} \quad f(\mathbf{x}) \\ & \text{subject to} \quad \mathbf{x} \in \mathbb{R}^n \end{aligned}$$

$\mathbf{x}^* \in \mathbb{R}^n$ を問題 P1 の大域的最適解とすると, 以下の不等式が成り立つことを示せ.

$$(\mathbf{x}^*)^\top \mathbf{x}^* \leq \frac{\mathbf{b}^\top \mathbf{b}}{\lambda_{\min}(\mathbf{C})}$$

ただし, $\lambda_{\min}(\mathbf{C})$ は \mathbf{C} の最小固有値を表す.

- (iv) \mathbf{A} を $m \times n$ 零行列, \mathbf{b} を m 次元零ベクトルと仮定する. 以下の最適化問題を考える.

$$\begin{aligned} \text{P2:} \quad & \text{Minimize} \quad f(\mathbf{x}) \\ & \text{subject to} \quad \mathbf{x}^\top \mathbf{x} \leq \alpha \end{aligned}$$

ここで, $\alpha \in \mathbb{R}$ は正の定数である. $(\hat{\mathbf{x}}, \rho), (\bar{\mathbf{x}}, \rho) \in \mathbb{R}^n \times \mathbb{R}$ が共に問題 P2 のカルーシュ・キューン・タッカー条件を満たすとき, $f(\hat{\mathbf{x}}) = f(\bar{\mathbf{x}})$ が成り立つことを示せ.

An English Translation:

Operations Research

3

Let $\mathbf{A} \in \mathbb{R}^{m \times n}$, $\mathbf{b} \in \mathbb{R}^m$ and $\mathbf{C} \in \mathbb{R}^{n \times n}$. Consider the following nonlinear programming problem with parameter $\mathbf{x} = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$:

$$\begin{aligned} \text{P}(\mathbf{x}): \quad & \text{Minimize} \quad \sum_{i=1}^n (\mathbf{z}^i)^\top \mathbf{z}^i + \mathbf{y}^\top \mathbf{y} + \mathbf{x}^\top \mathbf{C} \mathbf{x} \\ & \text{subject to} \quad \mathbf{y} - \sum_{i=1}^n x_i \mathbf{z}^i = \mathbf{A} \mathbf{x} - \mathbf{b}, \end{aligned}$$

where the decision variables are $\mathbf{y}, \mathbf{z}^i \in \mathbb{R}^m$ ($i = 1, \dots, n$), with $^\top$ denoting transposition. Moreover, denote by $f(\mathbf{x})$ the optimal value of problem $\text{P}(\mathbf{x})$, assuming that it is well-defined for all \mathbf{x} .

Answer the following questions.

- (i) Write out the Karush-Kuhn-Tucker conditions of $\text{P}(\mathbf{x})$.
- (ii) Prove that the objective function of problem $\text{P}(\mathbf{x})$ is convex with respect to $\mathbf{y}, \mathbf{z}^i \in \mathbb{R}^m$ ($i = 1, \dots, n$).
- (iii) Assume that \mathbf{C} is symmetric positive definite and consider the following optimization problem:

$$\begin{aligned} \text{P1:} \quad & \text{Minimize} \quad f(\mathbf{x}) \\ & \text{subject to} \quad \mathbf{x} \in \mathbb{R}^n. \end{aligned}$$

Show that the following inequality holds when $\mathbf{x}^* \in \mathbb{R}^n$ is a global optimal solution of problem P1:

$$(\mathbf{x}^*)^\top \mathbf{x}^* \leq \frac{\mathbf{b}^\top \mathbf{b}}{\lambda_{\min}(\mathbf{C})},$$

where $\lambda_{\min}(\mathbf{C})$ denotes the smallest eigenvalue of \mathbf{C} .

- (iv) Assume that \mathbf{A} is the $m \times n$ zero matrix and \mathbf{b} is the m -dimensional zero vector. Consider the following optimization problem:

$$\begin{aligned} \text{P2:} \quad & \text{Minimize} \quad f(\mathbf{x}) \\ & \text{subject to} \quad \mathbf{x}^\top \mathbf{x} \leq \alpha, \end{aligned}$$

where $\alpha \in \mathbb{R}$ is a positive constant. Show that $f(\hat{\mathbf{x}}) = f(\bar{\mathbf{x}})$ holds, when both $(\hat{\mathbf{x}}, \rho), (\bar{\mathbf{x}}, \rho) \in \mathbb{R}^n \times \mathbb{R}$ satisfy the Karush-Kuhn-Tucker conditions of problem P2.